



Delivering a Healthy WA

West Australian Vaccination Schedule

1st April 2008

BIRTH TO 4 YEARS	NON - INDIGENOUS VACCINE ANTIGENS	INDIGENOUS VACCINE ANTIGENS	BRAND NAME
Birth	Hep B (Paed)	Hep B (Paed)	H-B-VAX II (PAED)
2 months	7vPCV DTPa-IPV-HepB-Hib Rotavirus (ORV)*	7vPCV DTPa-IPV-HepB Hib Rotavirus (ORV)*	PREVENAR INFANRIX HEXA INFANRIX PENTA PEDVAXHIB ROTARIX
4 months	7vPCV DTPa-IPV-HepB-Hib Rotavirus (ORV)	7vPCV DTPa-IPV-HepB Hib Rotavirus (ORV)	PREVENAR INFANRIX HEXA INFANRIX PENTA PEDVAXHIB ROTARIX
6 months	7vPCV DTPa-IPV-HepB-Hib	7vPCV DTPa-IPV-HepB	PREVENAR INFANRIX HEXA INFANRIX PENTA
12 months	MMR Hib MenCCV	MMR Hib MenCCV Hep A	PRIORIX HIBERIX NEISVAC-C VAQTA
18 months	VZV	VZV 23vPPV Hep A	VARILRIX PNEUMOVAX23 VAQTA
4 years	MMR DTPa-IPV	MMR DTPa-IPV	PRIORIX INFANRIX-IPV

ADOLESCENT and ADULT	NON INDIGENOUS AND INDIGENOUS VACCINE ANTIGENS	BRAND NAME
Year 7 (primary school) females	Hep B (two Adult doses - 4-6 months apart) dTpa VZV HPV (1 dose given in terms 1,2 and 4)	H-B-VAX II (ADULT) BOOSTRIX VARILRIX GARDASIL
<u>Non-Indigenous</u> - 65 years or older <u>Indigenous</u> - 50 years or older <u>OR</u> 15-49 yrs (risk group)	Influenza (one dose - every year) 23vPPV (two doses - 5 or more years apart) <i>Indigenous 15-49 yrs (risk group) require an additional 23vPPV booster dose at 50 years of age OR 10 years after the 2nd dose, whichever is later</i>	FLUVAX, VAXIGRIP PNEUMOVAX23

Hib: *Indigenous children* - Primary schedule of 2 doses of PRP-OMP Hib vaccine (PedvaxHib) at 2 mths and 4 mths and a booster dose PRP-T-Hib (Hiberix) at 12 mths. If any dose of PRP-T-Hib vaccine (Infanrix Hexa or Hiberix) is administered at 2 or 4 months, a total of 3 doses of a Hib containing vaccine should be given in the primary schedule, with a booster dose of Hiberix at 12 months. *Non-Indigenous children* - Primary schedule of 3 doses of PRP-T-Hib vaccine (Infanrix Hexa) at 2 mths, 4 mths, 6 mths with a booster dose of Hiberix at 12 mths.

Hepatitis B: Give a total of three doses of Hepatitis B containing vaccine after the birth dose at either 2 mths, 4 mths and 6 mths; or at 2 mths, 4 mths and 12 mths.

Medical at-risk children: Require a 4th dose of 7vPCV at 12 mths of age, and a booster dose of 23vPPV at 4 yrs of age.

*** Rotavirus vaccine (Rotarix-ORV):** The Rotarix vaccine is for ORAL use. Two dose course, first dose between 6 and 14 weeks of age, second between 10 and 24 weeks of age. Minimum interval of 4 weeks. Late or catch up doses are not to be given.

Gardasil 12-26 years females HPV catch up 2008: School Years 8,9,10 (1 dose given in terms 1,2 and 4). 18 - 26 years & those missed out at school (GP based) 3 doses with recommended scheduled dose intervals of 0, 2 and 6 months.

