

Appendix 1

Site specific tables of incidence and mortality by age and sex

List of site specific tables and ICD codes

Table A1 –	All cancers (excluding NMSC) (140-208)
Table A2 –	Cancer of the Lip (140)
Table A3 –	Cancer of the Lip, & skin of lip (non-melanoma) (140, 173.0)
Table A4 –	Cancer of the Tongue (141)
Table A5 –	Cancer of the Salivary glands (142)
Table A6 –	Cancer of the Gum (143)
Table A7 –	Cancer of the Floor of mouth (144)
Table A8 –	Cancer of other & unspecified parts of the mouth (145)
Table A9 –	Cancer of the Buccal cavity (140-145, 173.0)
Table A10 –	Cancer of the Oropharynx (146)
Table A11 –	Cancer of the Nasopharynx (147)
Table A12 –	Cancer of the Hypopharynx (148)
Table A13 –	Cancers of ill-defined sites within the lip, oral cavity & pharynx (149)
Table A14 –	Cancer of the Oesophagus (150)
Table A15 –	Cancer of the Stomach (151)
Table A16 –	Cancer of the Small intestine, incl. duodenum (152)
Table A17 –	Cancer of the Large intestine, excluding rectum (153)
Table A18 –	Cancer of the Hepatic flexure (153.0)
Table A19 –	Cancer of the Transverse colon (153.1)
Table A20 –	Cancer of the Descending colon (153.2)
Table A21 –	Cancer of the Sigmoid colon (153.3)
Table A22 –	Cancer of the Caecum (153.4)
Table A23 –	Cancer of the Appendix (153.5)
Table A24 –	Cancer of the Ascending colon (153.6)
Table A25 –	Cancer of the Splenic flexure (153.7)
Table A26 –	Cancer of other specified sites of the Colon (153.8)
Table A27 –	Cancer of the Colon, unspecified (153.9)
Table A28 –	Cancer of the Rectum, rectosigmoid junction and anal canal (154)
Table A29 –	Cancer of the Rectosigmoid junction (154.0)
Table A30 –	Cancer of the Rectum (154.1)
Table A31 –	Cancer of the Anal canal (154.2)
Table A32 –	Cancer of the Anorectal junction (154.8)
Table A33 –	Cancer of the Colon and Rectum (including anus) (153-154)
Table A34 –	Cancer of the Liver & intrahepatic bile ducts (primary) (155)
Table A35 –	Cancer of the Gallbladder & extrahepatic bile ducts (156)
Table A36 –	Cancer of the Pancreas (157)
Table A37 –	Cancer of ill-defined sites - digestive organs (159)
Table A38 –	Cancer of the Digestive organs (150-159)
Table A39 –	Cancer of the Nose, nasal cavities, middle ear & sinuses (160)
Table A40 –	Cancer of the Larynx (161)
Table A41 –	Cancer of the Trachea, bronchus & lung (162)
Table A42 –	Cancer of the Pleura (163)
Table A43 –	Cancer of the Respiratory organs (160-163)
Table A44 –	Cancer of the Thymus, heart and mediastinum (164)
Table A45 –	Cancer of the Bone and articular cartilage (170)
Table A46 –	Cancer of the Connective & other soft tissue (171)
Table A47 –	Melanoma of skin (172)
Table A48 –	Melanoma of lip (172.0)
Table A49 –	Melanoma of eyelid, including canthus (172.1)
Table A50 –	Melanoma of ear (172.2)
Table A51 –	Melanoma of unspecified parts of face (172.3)
Table A52 –	Melanoma of scalp and neck (172.4)
Table A53 –	Melanoma of trunk (172.5)

Table A54 –	Melanoma of upper limb (172.6)
Table A55 –	Melanoma of lower limb (172.7)
Table A56 –	Melanoma of other site (172.8)
Table A57 –	Melanoma - site unspecified (172.9)
Table A58 –	Cancer of other skin - lip & anus (173)
Table A59 –	Cancer of the Skin of lip (173.0)
Table A60 –	Cancer of the Skin of anus (173.5)
Table A61 –	Cancer of the Female breast (174)
Table A62 –	Cancer of the Male breast (175)
Table A63 –	Cancer of the Cervix uteri, unspecified (180.9)
Table A64 –	Cancer of the Placenta (181)
Table A65 –	Cancer of the Body of the uterus (182)
Table A66 –	Cancer of the Ovary and other uterine adnexa (183)
Table A67 –	Cancer of the Other & unspecified female genitalia (184)
Table A68 –	Cancer of the Vagina & unspecified female genital organs (184.0 184.8 184.9)
Table A69 –	Cancer of the Vulva (184.1 184.2 184.3 184.4)
Table A70 –	Cancer of the Prostate (185)
Table A71 –	Cancer of the Testis (186)
Table A72 –	Cancer of the Penis and other male genital organs (187)
Table A73 –	Cancer of the Bladder (188)
Table A74 –	Cancer of the Kidney, and other unspec. urinary organs (189)
Table A75 –	Cancer of the Eye (190)
Table A76 –	Cancer of the Brain (191)
Table A77 –	Cancer of the Cranial nerves, spinal cord and meninges (192)
Table A78 –	Cancer of the Thyroid gland (193)
Table A79 –	Cancer of the other endocrine glands (194)
Table A80 –	Cancer of Unspecified site (199)
Table A81 –	Diffuse non-Hodgkin's lymphomas (200)
Table A82 –	Hodgkin's disease (201)
Table A83 –	Other lymphomas & histiocytic tissue tumours (202)
Table A84 –	Nodular lymphomas (202.0)
Table A85 –	All lymphomas (200 201 202.0 202.1 202.2 202.8. 202.9)
Table A86 –	Other lymphomas (202.1 202.2 202.8 202.9)
Table A87 –	Tumors of histiocytic tissue (202.3 202.5 202.6)
Table A88 –	Multiple myeloma & immunoproliferative neoplasms (203)
Table A89 –	All lymphoid leukaemias (204)
Table A90 –	Acute lymphatic leukaemia (204.0)
Table A91 –	Chronic lymphatic leukaemia (204.1)
Table A92 –	All myeloid leukaemias (205)
Table A93 –	Acute myeloid leukaemia (205.0)
Table A94 –	Chronic myeloid leukaemia (205.0)
Table A95 –	Chronic monocytic leukaemia (205.0)
Table A96 –	All leukaemias (204 205 206.1 207 208)
Table A97 –	Monocytic leukaemias (chronic only) (206)
Table A98 –	Other specified leukaemias (207)
Table A99 –	Unspecified cell leukaemias (208)
Table A100 –	All cancers (140–208)
Table A101 –	Cancer of the Head and Neck (141–149)

Table A1 – All cancers (excluding NMSC) (140–208)

South Australia Year – 2002	Incidence						Mortality					
	New cases			Rates			Deaths			Rates		
	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons
0–4	9	6	15	19.4	13.5	16.5	2	0	2	4.3	0.0	2.2
5–9	7	7	14	13.9	14.6	14.2	1	2	3	2.0	4.2	3.1
10–14	6	6	12	11.6	12.3	11.9	2	1	3	3.9	2.0	3.0
15–19	22	12	34	41.6	23.8	32.9	3	3	6	5.7	5.9	5.8
20–24	16	17	33	32.2	36.1	34.1	1	1	2	2.0	2.1	2.1
25–29	29	32	61	58.6	67.4	62.9	5	4	9	10.1	8.4	9.3
30–34	44	68	112	79.8	126.0	102.6	10	5	15	18.1	9.3	13.7
35–39	74	81	155	132.3	146.1	139.1	15	19	34	26.8	34.3	30.5
40–44	99	154	253	169.9	261.5	215.9	17	31	48	29.2	52.6	41.0
45–49	147	207	354	273.9	378.3	326.6	33	54	87	61.5	98.7	80.3
50–54	244	343	587	472.8	649.9	562.3	83	96	179	160.8	181.9	171.5
55–59	350	353	703	793.6	794.7	794.1	131	99	230	297.0	222.9	259.8
60–64	431	353	784	1,264.6	1,016.9	1,139.6	143	110	253	419.6	316.9	367.8
65–69	527	382	909	1,828.8	1,261.2	1,538.0	219	150	369	760.0	495.2	624.3
70–74	735	411	1,146	2,767.6	1,382.7	2,036.2	288	173	461	1,084.5	582.0	819.1
75–79	652	436	1,088	3,020.8	1,566.0	2,201.3	356	229	585	1,649.4	822.5	1,183.6
80–84	410	363	773	3,190.7	1,803.7	2,344.2	269	235	504	2,093.4	1,167.7	1,528.4
85+	289	386	675	3,602.1	2,111.7	2,566.3	243	291	534	3,028.8	1,592.0	2,030.3
Total/Crude rate	4,091	3,617	7,708	544.5	471.3	507.5	1,821	1,503	3,324	242.4	195.9	218.9
AS Rate (Aust)				526.7	398.0	451.9				239.4	153.2	188.7
Confidence Interval (95%)				510.5 – 542.9	384.8 – 411.2	441.7 – 462.0				228.3 – 250.4	145.3 – 161.1	182.3 – 195.2
AS Rate (World)				378.2	303.8	335.5				157.5	106.3	128.1
Confidence Interval (95%)				367.3 – 389.1	293.9 – 313.7	328.2 – 342.8				151.0 – 164.1	100.9 – 111.7	124.0 – 132.3
Risk				1 in 3	1 in 4	1 in 3				1 in 7	1 in 10	1 in 9
PYLL										11,558	10,855	22,413
% of all cancers	100.0	100.0	100.0				100.0	100.0	100.0	100.0	100.0	100.0
Trend 1998–2002 (per annum)				-1.0%	-0.3%	-0.5%				-0.8%	0.3%	-0.3%

Table A2 – Cancer of the Lip (140)

South Australia Year – 2002	Incidence						Mortality					
	New cases			Rates			Deaths			Rates		
	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons
0–4	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
5–9	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
10–14	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
15–19	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
20–24	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
25–29	3	0	3	6.1	0.0	3.1	0	0	0	0.0	0.0	0.0
30–34	5	1	6	9.1	1.9	5.5	0	0	0	0.0	0.0	0.0
35–39	4	2	6	7.2	3.6	5.4	0	0	0	0.0	0.0	0.0
40–44	9	1	10	15.4	1.7	8.5	0	0	0	0.0	0.0	0.0
45–49	9	3	12	16.8	5.5	11.1	0	0	0	0.0	0.0	0.0
50–54	8	3	11	15.5	5.7	10.5	0	0	0	0.0	0.0	0.0
55–59	11	1	12	24.9	2.3	13.6	1	0	1	2.3	0.0	1.1
60–64	9	3	12	26.4	8.6	17.4	0	0	0	0.0	0.0	0.0
65–69	6	2	8	20.8	6.6	13.5	0	0	0	0.0	0.0	0.0
70–74	9	3	12	33.9	10.1	21.3	0	0	0	0.0	0.0	0.0
75–79	8	6	14	37.1	21.6	28.3	0	0	0	0.0	0.0	0.0
80–84	8	6	14	62.3	29.8	42.5	0	0	0	0.0	0.0	0.0
85+	5	8	13	62.3	43.8	49.4	0	0	0	0.0	0.0	0.0
Total/Crude rate	94	39	133	12.5	5.1	8.8	1	0	1	0.1	0.0	0.1
AS Rate (Aust)				12.2	4.0	8.0				0.1	0.0	0.1
Confidence Interval (95%)				9.7 – 14.7	2.7 – 5.3	6.6 – 9.3				0.0 – 0.3	0.0 – 0.0	0.0 – 0.2
AS Rate (World)				9.5	2.8	6.1				0.1	0.0	0.1
Confidence Interval (95%)				7.7 – 11.3	2.0 – 3.7	5.1 – 7.1				0.0 – 0.3	0.0 – 0.0	0.0 – 0.1
Risk				1 in 114	1 in 436	1 in 182				1 in 8,821	N/A	1 in 17,705
PYLL										18	0	18
% of all cancers	2.3	1.1	1.7				2.3	1.1	1.7	0.1	0.0	0.0
Trend 1998–2002 (per annum)				2.1%	1.5%	2.4%				-6.1%	-17.0%	-4.9%

Table A3 – Cancer of the Lip, & skin of lip (non-melanoma) (140, 173.0)

South Australia Year – 2002	Incidence						Mortality					
	New cases			Rates			Deaths			Rates		
	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons
0–4	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
5–9	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
10–14	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
15–19	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
20–24	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
25–29	3	0	3	6.1	0.0	3.1	0	0	0	0.0	0.0	0.0
30–34	7	1	8	12.7	1.9	7.3	0	0	0	0.0	0.0	0.0
35–39	5	3	8	8.9	5.4	7.2	0	0	0	0.0	0.0	0.0
40–44	12	1	13	20.6	1.7	11.1	0	0	0	0.0	0.0	0.0
45–49	11	4	15	20.5	7.3	13.8	0	0	0	0.0	0.0	0.0
50–54	12	4	16	23.3	7.6	15.3	0	0	0	0.0	0.0	0.0
55–59	13	5	18	29.5	11.3	20.3	1	0	1	2.3	0.0	1.1
60–64	15	4	19	44.0	11.5	27.6	0	0	0	0.0	0.0	0.0
65–69	9	5	14	31.2	16.5	23.7	0	0	0	0.0	0.0	0.0
70–74	11	5	16	41.4	16.8	28.4	0	0	0	0.0	0.0	0.0
75–79	11	7	18	51.0	25.1	36.4	1	0	1	4.6	0.0	2.0
80–84	9	8	17	70.0	39.8	51.6	0	0	0	0.0	0.0	0.0
85+	7	13	20	87.2	71.1	76.0	0	1	1	0.0	5.5	3.8
Total/Crude rate	125	60	185	16.6	7.8	12.2	2	1	3	0.3	0.1	0.2
AS Rate (Aust)				16.2	6.2	11.1				0.2	0.1	0.2
Confidence Interval (95%)				13.4 – 19.1	4.6 – 7.8	9.5 – 12.7				0.0 – 0.6	0.0 – 0.2	0.0 – 0.4
AS Rate (World)				12.7	4.5	8.5				0.2	0.0	0.1
Confidence Interval (95%)				10.6 – 14.8	3.3 – 5.6	7.3 – 9.6				0.0 – 0.4	0.0 – 0.1	0.0 – 0.2
Risk				1 in 84	1 in 251	1 in 127				1 in 8,821	N/A	1 in 17,705
PYLL										18	0	18
% of all cancers	3.1	1.7	2.4				3.1	1.7	2.4	0.1	0.1	0.1
Trend 1998–2002 (per annum)				-3.5%	-4.6%	-3.6%				-1.1%	-3.9%	4.4%

Note: All rates are expressed per 100,000 population. Age-standardised rates (AS Rate) use the Australian 2001 and New World Standard populations (see Methods)

Table A4 – Cancer of the Tongue (141)

South Australia Year – 2002	Incidence						Mortality					
	New cases			Rates			Deaths			Rates		
	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons
0–4	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
5–9	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
10–14	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
15–19	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
20–24	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
25–29	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
30–34	1	0	1	1.8	0.0	0.9	0	0	0	0.0	0.0	0.0
35–39	2	0	2	3.6	0.0	1.8	0	0	0	0.0	0.0	0.0
40–44	2	0	2	3.4	0.0	1.7	0	0	0	0.0	0.0	0.0
45–49	1	1	2	1.9	1.8	1.8	0	0	0	0.0	0.0	0.0
50–54	5	2	7	9.7	3.8	6.7	1	0	1	1.9	0.0	1.0
55–59	0	2	2	0.0	4.5	2.3	2	0	2	4.5	0.0	2.3
60–64	3	0	3	8.8	0.0	4.4	1	0	1	2.9	0.0	1.5
65–69	3	0	3	10.4	0.0	5.1	0	0	0	0.0	0.0	0.0
70–74	3	2	5	11.3	6.7	8.9	2	1	3	7.5	3.4	5.3
75–79	2	2	4	9.3	7.2	8.1	1	0	1	4.6	0.0	2.0
80–84	0	0	0	0.0	0.0	0.0	1	0	1	7.8	0.0	3.0
85+	0	1	1	0.0	5.5	3.8	0	1	1	0.0	5.5	3.8
Total/Crude rate	22	10	32	2.9	1.3	2.1	8	2	10	1.1	0.3	0.7
AS Rate (Aust)				2.8	1.1	2.0				1.0	0.2	0.6
Confidence Interval (95%)				1.6–4.0	0.4–1.8	1.3–2.6				0.3–1.7	0.0–0.4	0.2–0.9
AS Rate (World)				2.3	0.8	1.5				0.7	0.1	0.4
Confidence Interval (95%)				1.4–3.2	0.3–1.3	1.0–2.0				0.3–1.2	0.0–0.2	0.2–0.6
Risk				1 in 394	1 in 1,188	1 in 597				1 in 1,181	1 in 5,945	1 in 2,000
PYLL										75	3	78
% of all cancers	0.5	0.3	0.4				0.5	0.3	0.4	0.4	0.1	0.3
Trend 1998–2002 (per annum)				16.3%	-7.8%	5.4%				-6.5%	-12.0%	-8.2%

Table A5 – Cancer of the Salivary glands (142)

South Australia Year – 2002	Incidence						Mortality					
	New cases			Rates			Deaths			Rates		
	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons
0–4	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
5–9	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
10–14	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
15–19	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
20–24	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
25–29	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
30–34	0	1	1	0.0	1.9	0.9	0	0	0	0.0	0.0	0.0
35–39	1	1	2	1.8	1.8	1.8	0	0	0	0.0	0.0	0.0
40–44	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
45–49	1	1	2	1.9	1.8	1.8	0	0	0	0.0	0.0	0.0
50–54	2	1	3	3.9	1.9	2.9	2	0	2	3.9	0.0	1.9
55–59	1	0	1	2.3	0.0	1.1	1	0	1	2.3	0.0	1.1
60–64	1	1	2	2.9	2.9	2.9	1	0	1	2.9	0.0	1.5
65–69	1	0	1	3.5	0.0	1.7	1	0	1	3.5	0.0	1.7
70–74	5	1	6	18.8	3.4	10.7	0	0	0	0.0	0.0	0.0
75–79	1	2	3	4.6	7.2	6.1	1	0	1	4.6	0.0	2.0
80–84	2	1	3	15.6	5.0	9.1	1	2	3	7.8	9.9	9.1
85+	1	1	2	12.5	5.5	7.6	0	2	2	0.0	10.9	7.6
Total/Crude rate	16	10	26	2.1	1.3	1.7	7	4	11	0.9	0.5	0.7
AS Rate (Aust)				2.1	1.1	1.5				0.9	0.3	0.6
Confidence Interval (95%)				1.1–3.1	0.4–1.8	0.9–2.1				0.2–1.5	0.0–0.6	0.3–1.0
AS Rate (World)				1.5	0.8	1.1				0.7	0.2	0.4
Confidence Interval (95%)				0.8–2.1	0.3–1.4	0.7–1.5				0.2–1.1	0.1–0.3	0.2–0.7
Risk				1 in 572	1 in 1,469	1 in 840				1 in 1,594	N/A	1 in 3,231
PYLL										83	0	83
% of all cancers	0.4	0.3	0.3				0.4	0.3	0.3	0.4	0.3	0.3
Trend 1998–2002 (per annum)				1.2%	21.3%	9.0%				29.3%	11.1%	23.6%

Table A6 – Cancer of the Gum (143)

South Australia Year – 2002	Incidence						Mortality					
	New cases			Rates			Deaths			Rates		
	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons
0–4	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
5–9	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
10–14	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
15–19	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
20–24	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
25–29	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
30–34	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
35–39	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
40–44	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
45–49	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
50–54	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
55–59	1	0	1	2.3	0.0	1.1	0	0	0	0.0	0.0	0.0
60–64	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
65–69	1	0	1	3.5	0.0	1.7	1	0	1	3.5	0.0	1.7
70–74	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
75–79	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
80–84	0	0	0	0.0	0.0	0.0	1	0	1	7.8	0.0	3.0
85+	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
Total/Crude rate	2	0	2	0.3	0.0	0.1	2	0	2	0.3	0.0	0.1
AS Rate (Aust)				0.2	0.0	0.1				0.3	0.0	0.1
Confidence Interval (95%)				0.0–0.6	0.0–0.0	0.0–0.3				0.0–0.6	0.0–0.0	0.0–0.3
AS Rate (World)				0.2	0.0	0.1				0.2	0.0	0.1
Confidence Interval (95%)				0.0–0.5	0.0–0.0	0.0–0.2				0.0–0.4	0.0–0.0	0.0–0.2
Risk				1 in 3,486	N/A	1 in 7,089				1 in 5,764	N/A	1 in 11,821
PYLL										8	0	8
% of all cancers	0.0	0.0	0.0				0.0	0.0	0.0	0.1	0.0	0.1
Trend 1998–2002 (per annum)				11.7%	-24.9%	-12.2%				-18.2%	-30.7%	-23.0%

Note: All rates are expressed per 100,000 population. Age-standardised rates (AS Rate) use the Australian 2001 and New World Standard populations (see Methods)

Table A7 – Cancer of the Floor of mouth (144)

South Australia Year – 2002	Incidence						Mortality					
	New cases			Rates			Deaths			Rates		
	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons
0-4	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
5-9	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
10-14	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
15-19	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
20-24	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
25-29	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
30-34	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
35-39	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
40-44	0	0	0	0.0	0.0	0.0	1	1	2	1.7	1.7	1.7
45-49	0	1	1	0.0	1.8	0.9	1	0	1	1.9	0.0	0.9
50-54	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
55-59	1	0	1	2.3	0.0	1.1	0	0	0	0.0	0.0	0.0
60-64	1	0	1	2.9	0.0	1.5	2	0	2	5.9	0.0	2.9
65-69	2	1	3	6.9	3.3	5.1	0	0	0	0.0	0.0	0.0
70-74	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
75-79	2	1	3	9.3	3.6	6.1	0	0	0	0.0	0.0	0.0
80-84	1	0	1	7.8	0.0	3.0	0	0	0	0.0	0.0	0.0
85+	1	0	1	12.5	0.0	3.8	1	0	1	12.5	0.0	3.8
Total/Crude rate	8	3	11	1.1	0.4	0.7	5	1	6	0.7	0.1	0.4
AS Rate (Aust)				1.0	0.3	0.6				0.7	0.1	0.4
Confidence Interval (95%)				0.3 – 1.8	0.0 – 0.7	0.3 – 1.0				0.1 – 1.3	0.0 – 0.4	0.1 – 0.7
AS Rate (World)				0.7	0.3	0.5				0.5	0.1	0.3
Confidence Interval (95%)				0.3 – 1.2	0.0 – 0.6	0.2 – 0.7				0.1 – 1.0	0.0 – 0.3	0.1 – 0.5
Risk				1 in 1,648	1 in 3,900	1 in 2,331				1 in 2,117	1 in 11,777	1 in 3,613
PYLL										85	33	118
% of all cancers	0.2	0.1	0.1				0.2	0.1	0.1	0.3	0.1	0.2
Trend 1998-2002 (per annum)				-9.1%	-12.2%	-11.0%				5.0%	-19.6%	-6.9%

Table A8 – Cancer of other & unspecified parts of the mouth (145)

South Australia Year – 2002	Incidence						Mortality					
	New cases			Rates			Deaths			Rates		
	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons
0-4	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
5-9	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
10-14	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
15-19	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
20-24	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
25-29	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
30-34	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
35-39	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
40-44	1	0	1	1.7	0.0	0.9	0	0	0	0.0	0.0	0.0
45-49	0	1	1	0.0	1.8	0.9	0	0	0	0.0	0.0	0.0
50-54	2	1	3	3.9	1.9	2.9	0	0	0	0.0	0.0	0.0
55-59	1	1	2	2.3	2.3	2.3	0	0	0	0.0	0.0	0.0
60-64	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
65-69	1	1	2	3.5	3.3	3.4	0	0	0	0.0	0.0	0.0
70-74	0	2	2	0.0	6.7	3.6	0	0	0	0.0	0.0	0.0
75-79	3	1	4	13.9	3.6	8.1	0	0	0	0.0	0.0	0.0
80-84	0	1	1	0.0	5.0	3.0	1	0	1	7.8	0.0	3.0
85+	0	2	2	0.0	10.9	7.6	2	0	2	24.9	0.0	7.6
Total/Crude rate	8	10	18	1.1	1.3	1.2	3	0	3	0.4	0.0	0.2
AS Rate (Aust)				1.0	1.0	1.0				0.5	0.0	0.2
Confidence Interval (95%)				0.3 – 1.7	0.4 – 1.7	0.6 – 1.5				0.0 – 1.0	0.0 – 0.0	0.0 – 0.3
AS Rate (World)				0.7	0.7	0.7				0.2	0.0	0.1
Confidence Interval (95%)				0.3 – 1.2	0.3 – 1.2	0.4 – 1.1				0.0 – 0.4	0.0 – 0.0	0.0 – 0.1
Risk				1 in 1,766	1 in 1,250	1 in 1,445				N/A	N/A	N/A
PYLL										0	0	0
% of all cancers	0.2	0.3	0.2				0.2	0.3	0.2	0.2	0.0	0.1
Trend 1998-2002 (per annum)				-7.6%	-4.3%	-6.6%				10.6%	-19.1%	-8.3%

Table A9 – Cancer of the Buccal cavity (140-145, 173.0)

South Australia Year – 2002	Incidence						Mortality					
	New cases			Rates			Deaths			Rates		
	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons
0-4	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
5-9	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
10-14	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
15-19	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
20-24	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
25-29	3	0	3	6.1	0.0	3.1	0	0	0	0.0	0.0	0.0
30-34	8	2	10	14.5	3.7	9.2	0	0	0	0.0	0.0	0.0
35-39	8	4	12	14.3	7.2	10.8	0	0	0	0.0	0.0	0.0
40-44	15	1	16	25.7	1.7	13.7	1	1	2	1.7	1.7	1.7
45-49	13	8	21	24.2	14.6	19.4	1	0	1	1.9	0.0	0.9
50-54	21	8	29	40.7	15.2	27.8	3	0	3	5.8	0.0	2.9
55-59	17	8	25	38.5	18.0	28.2	4	0	4	9.1	0.0	4.5
60-64	20	5	25	58.7	14.4	36.3	4	0	4	11.7	0.0	5.8
65-69	17	7	24	59.0	23.1	40.6	2	0	2	6.9	0.0	3.4
70-74	19	10	29	71.5	33.6	51.5	2	1	3	7.5	3.4	5.3
75-79	19	13	32	88.0	46.7	64.7	3	0	3	13.9	0.0	6.1
80-84	12	10	22	93.4	49.7	66.7	4	2	6	31.1	9.9	18.2
85+	9	17	26	112.2	93.0	98.9	3	4	7	37.4	21.9	26.6
Total/Crude rate	181	93	274	24.1	12.1	18.0	27	8	35	3.6	1.0	2.3
AS Rate (Aust)				23.4	9.8	16.3				3.5	0.7	2.0
Confidence Interval (95%)				20.0 – 26.8	7.8 – 11.9	14.4 – 18.3				2.2 – 4.9	0.2 – 1.2	1.3 – 2.7
AS Rate (World)				18.1	7.1	12.4				2.5	0.4	1.4
Confidence Interval (95%)				15.6 – 20.6	5.7 – 8.5	11.0 – 13.9				1.6 – 3.3	0.1 – 0.7	0.9 – 1.8
Risk				1 in 57	1 in 153	1 in 84				1 in 448	1 in 3,951	1 in 815
PYLL										268	35	303
% of all cancers	4.4	2.6	3.6				4.4	2.6	3.6	1.5	0.5	1.1
Trend 1998-2002 (per annum)				-2.2%	-4.3%	-2.7%				1.9%	-12.7%	-4.5%

Note: All rates are expressed per 100,000 population. Age-standardised rates (AS Rate) use the Australian 2001 and New World Standard populations (see Methods)

Table A10 – Cancer of the Oropharynx (146)

South Australia Year – 2002	Incidence						Mortality					
	New cases			Rates			Deaths			Rates		
	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons
0–4	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
5–9	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
10–14	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
15–19	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
20–24	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
25–29	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
30–34	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
35–39	2	0	2	3.6	0.0	1.8	0	0	0	0.0	0.0	0.0
40–44	3	0	3	5.1	0.0	2.6	0	0	0	0.0	0.0	0.0
45–49	5	1	6	9.3	1.8	5.5	3	0	3	5.6	0.0	2.8
50–54	3	0	3	5.8	0.0	2.9	1	0	1	1.9	0.0	1.0
55–59	2	0	2	4.5	0.0	2.3	1	0	1	2.3	0.0	1.1
60–64	4	0	4	11.7	0.0	5.8	1	1	2	2.9	2.9	2.9
65–69	0	0	0	0.0	0.0	0.0	1	0	1	3.5	0.0	1.7
70–74	4	1	5	15.1	3.4	8.9	1	0	1	3.8	0.0	1.8
75–79	3	1	4	13.9	3.6	8.1	1	0	1	4.6	0.0	2.0
80–84	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
85+	0	0	0	0.0	0.0	0.0	1	0	1	12.5	0.0	3.8
Total/Crude rate	26	3	29	3.5	0.4	1.9	10	1	11	1.3	0.1	0.7
AS Rate (Aust)				3.3	0.3	1.8				1.3	0.1	0.7
Confidence Interval (95%)				2.0 – 4.6	0.0 – 0.7	1.1 – 2.4				0.5 – 2.1	0.0 – 0.4	0.3 – 1.1
AS Rate (World)				2.7	0.2	1.4				1.0	0.1	0.5
Confidence Interval (95%)				1.7 – 3.6	0.0 – 0.5	0.9 – 1.9				0.4 – 1.6	0.0 – 0.3	0.2 – 0.8
Risk				1 in 363	1 in 3,853	1 in 673				1 in 1,002	1 in 6,943	1 in 1,781
PYLL										145	13	158
% of all cancers	0.6	0.1	0.4				0.6	0.1	0.4	0.5	0.1	0.3
Trend 1998–2002 (per annum)				19.5%	-14.2%	8.0%				-3.1%	-17.6%	-7.6%

Table A11 – Cancer of the Nasopharynx (147)

South Australia Year – 2002	Incidence						Mortality					
	New cases			Rates			Deaths			Rates		
	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons
0–4	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
5–9	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
10–14	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
15–19	1	0	1	1.9	0.0	1.0	0	0	0	0.0	0.0	0.0
20–24	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
25–29	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
30–34	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
35–39	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
40–44	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
45–49	0	0	0	0.0	0.0	0.0	0	1	1	0.0	1.8	0.9
50–54	1	0	1	1.9	0.0	1.0	0	0	0	0.0	0.0	0.0
55–59	2	1	3	4.5	2.3	3.4	1	0	1	2.3	0.0	1.1
60–64	1	0	1	2.9	0.0	1.5	0	0	0	0.0	0.0	0.0
65–69	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
70–74	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
75–79	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
80–84	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
85+	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
Total/Crude rate	5	1	6	0.7	0.1	0.4	1	1	2	0.1	0.1	0.1
AS Rate (Aust)				0.6	0.1	0.4				0.1	0.1	0.1
Confidence Interval (95%)				0.1 – 1.2	0.0 – 0.3	0.1 – 0.7				0.0 – 0.3	0.0 – 0.4	0.0 – 0.3
AS Rate (World)				0.6	0.1	0.3				0.1	0.1	0.1
Confidence Interval (95%)				0.1 – 1.1	0.0 – 0.3	0.1 – 0.6				0.0 – 0.3	0.0 – 0.3	0.0 – 0.2
Risk				1 in 1,771	1 in 8,885	1 in 2,956				1 in 8,821	1 in 10,944	1 in 9,746
PYLL										18	28	45
% of all cancers	0.1	0.0	0.1				0.1	0.0	0.1	0.1	0.1	0.1
Trend 1998–2002 (per annum)				5.7%	-9.9%	2.0%				25.3%	27.3%	26.0%

Table A12 – Cancer of the Hypopharynx (148)

South Australia Year – 2002	Incidence						Mortality					
	New cases			Rates			Deaths			Rates		
	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons
0–4	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
5–9	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
10–14	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
15–19	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
20–24	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
25–29	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
30–34	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
35–39	2	0	2	3.6	0.0	1.8	0	0	0	0.0	0.0	0.0
40–44	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
45–49	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
50–54	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
55–59	1	0	1	2.3	0.0	1.1	1	0	1	2.3	0.0	1.1
60–64	1	0	1	2.9	0.0	1.5	2	0	2	5.9	0.0	2.9
65–69	1	0	1	3.5	0.0	1.7	0	0	0	0.0	0.0	0.0
70–74	1	0	1	3.8	0.0	1.8	0	0	0	0.0	0.0	0.0
75–79	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
80–84	1	0	1	7.8	0.0	3.0	1	0	1	7.8	0.0	3.0
85+	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
Total/Crude rate	7	0	7	0.9	0.0	0.5	4	0	4	0.5	0.0	0.3
AS Rate (Aust)				0.9	0.0	0.4				0.5	0.0	0.2
Confidence Interval (95%)				0.2 – 1.6	0.0 – 0.0	0.1 – 0.7				0.0 – 1.0	0.0 – 0.0	0.0 – 0.5
AS Rate (World)				0.7	0.0	0.4				0.4	0.0	0.2
Confidence Interval (95%)				0.2 – 1.2	0.0 – 0.0	0.1 – 0.6				0.0 – 0.8	0.0 – 0.0	0.0 – 0.4
Risk				1 in 1,250	N/A	1 in 2,549				1 in 2,459	N/A	1 in 4,955
PYLL										43	0	43
% of all cancers	0.2	0.0	0.1				0.2	0.0	0.1	0.2	0.0	0.1
Trend 1998–2002 (per annum)				-6.4%	-17.1%	-7.0%				-12.2%	8.7%	-9.3%

Note: All rates are expressed per 100,000 population. Age-standardised rates (AS Rate) use the Australian 2001 and New World Standard populations (see Methods)

Table A13 – Cancer off ill-defined sites within the lip, oral cavity & pharynx (149)

South Australia Year – 2002	Incidence						Mortality					
	New cases			Rates			Deaths			Rates		
	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons
0-4	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
5-9	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
10-14	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
15-19	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
20-24	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
25-29	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
30-34	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
35-39	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
40-44	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
45-49	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
50-54	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
55-59	1	0	1	2.3	0.0	1.1	0	0	0	0.0	0.0	0.0
60-64	1	0	1	2.9	0.0	1.5	0	0	0	0.0	0.0	0.0
65-69	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
70-74	1	0	1	3.8	0.0	1.8	1	0	1	3.8	0.0	1.8
75-79	0	0	0	0.0	0.0	0.0	1	1	2	4.6	3.6	4.0
80-84	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
85+	0	1	1	0.0	5.5	3.8	0	1	1	0.0	5.5	3.8
Total/Crude rate	3	1	4	0.4	0.1	0.3	2	2	4	0.3	0.3	0.3
AS Rate (Aust)				0.4	0.1	0.2				0.2	0.2	0.2
Confidence Interval (95%)				0.0 – 0.8	0.0 – 0.2	0.0 – 0.5				0.0 – 0.6	0.0 – 0.4	0.0 – 0.4
AS Rate (World)				0.3	0.0	0.2				0.2	0.1	0.1
Confidence Interval (95%)				0.0 – 0.6	0.0 – 0.1	0.0 – 0.3				0.0 – 0.3	0.0 – 0.2	0.0 – 0.2
Risk				1 in 2,231	N/A	1 in 4,588				1 in 5,312	N/A	1 in 11,257
PYLL										3	0	3
% of all cancers	0.1	0.0	0.1				0.1	0.0	0.1	0.1	0.1	0.1
Trend 1998-2002 (per annum)				58.1%	19.3%	46.4%				16.0%	29.5%	22.2%

Table A14 – Cancer of the Oesophagus (150)

South Australia Year – 2002	Incidence						Mortality					
	New cases			Rates			Deaths			Rates		
	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons
0-4	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
5-9	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
10-14	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
15-19	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
20-24	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
25-29	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
30-34	0	1	1	0.0	1.9	0.9	0	1	1	0.0	1.9	0.9
35-39	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
40-44	2	0	2	3.4	0.0	1.7	1	1	2	1.7	1.7	1.7
45-49	4	0	4	7.5	0.0	3.7	1	0	1	1.9	0.0	0.9
50-54	3	1	4	5.8	1.9	3.8	2	0	2	3.9	0.0	1.9
55-59	6	3	9	13.6	6.8	10.2	8	2	10	18.1	4.5	11.3
60-64	9	1	10	26.4	2.9	14.5	4	1	5	11.7	2.9	7.3
65-69	3	6	9	10.4	19.8	15.2	7	4	11	24.3	13.2	18.6
70-74	8	4	12	30.1	13.5	21.3	9	3	12	33.9	10.1	21.3
75-79	13	2	15	60.2	7.2	30.3	6	3	9	27.8	10.8	18.2
80-84	5	6	11	38.9	29.8	33.4	5	3	8	38.9	14.9	24.3
85+	3	5	8	37.4	27.4	30.4	8	5	13	99.7	27.4	49.4
Total/Crude rate	56	29	85	7.5	3.8	5.6	51	23	74	6.8	3.0	4.9
AS Rate (Aust)				7.1	3.0	4.9				6.7	2.3	4.2
Confidence Interval (95%)				5.3 – 9.0	1.9 – 4.0	3.8 – 5.9				4.8 – 8.6	1.4 – 3.3	3.3 – 5.2
AS Rate (World)				5.0	2.1	3.5				4.5	1.6	2.9
Confidence Interval (95%)				3.8 – 6.3	1.3 – 2.8	2.8 – 4.2				3.4 – 5.7	1.0 – 2.3	2.3 – 3.6
Risk				1 in 206	1 in 429	1 in 281				1 in 210	1 in 585	1 in 313
PYLL										370	160	530
% of all cancers	1.4	0.8	1.1				1.4	0.8	1.1	2.8	1.5	2.2
Trend 1998-2002 (per annum)				-4.7%	-4.4%	-4.6%				0.9%	0.1%	0.0%

Table A15 – Cancer of the Stomach (151)

South Australia Year – 2002	Incidence						Mortality					
	New cases			Rates			Deaths			Rates		
	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons
0-4	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
5-9	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
10-14	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
15-19	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
20-24	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
25-29	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
30-34	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
35-39	1	1	2	1.8	1.8	1.8	1	0	1	1.8	0.0	0.9
40-44	2	1	3	3.4	1.7	2.6	2	1	3	3.4	1.7	2.6
45-49	4	2	6	7.5	3.7	5.5	2	2	4	3.7	3.7	3.7
50-54	4	2	6	7.8	3.8	5.7	4	1	5	7.8	1.9	4.8
55-59	8	4	12	18.1	9.0	13.6	10	2	12	22.7	4.5	13.6
60-64	8	5	13	23.5	14.4	18.9	8	5	13	23.5	14.4	18.9
65-69	17	4	21	59.0	13.2	35.5	10	3	13	34.7	9.9	22.0
70-74	16	4	20	60.2	13.5	35.5	12	2	14	45.2	6.7	24.9
75-79	20	17	37	92.7	61.1	74.9	23	11	34	106.6	39.5	68.8
80-84	13	7	20	101.2	34.8	60.7	8	8	16	62.3	39.8	48.5
85+	12	10	22	149.6	54.7	83.6	11	12	23	137.1	65.6	87.4
Total/Crude rate	105	57	162	14.0	7.4	10.7	91	47	138	12.1	6.1	9.1
AS Rate (Aust)				13.7	5.7	9.2				11.8	4.6	7.8
Confidence Interval (95%)				11.0 – 16.3	4.2 – 7.2	7.8 – 10.6				9.4 – 14.3	3.2 – 5.9	6.5 – 9.1
AS Rate (World)				9.2	3.9	6.3				7.9	3.0	5.2
Confidence Interval (95%)				7.6 – 10.8	2.9 – 4.9	5.4 – 7.2				6.4 – 9.4	2.1 – 3.8	4.4 – 6.1
Risk				1 in 111	1 in 328	1 in 168				1 in 141	1 in 468	1 in 220
PYLL										628	235	863
% of all cancers	2.6	1.6	2.1				2.6	1.6	2.1	5.0	3.1	4.2
Trend 1998-2002 (per annum)				-4.1%	-4.6%	-4.3%				0.2%	-4.4%	-1.1%

Note: All rates are expressed per 100,000 population. Age-standardised rates (AS Rate) use the Australian 2001 and New World Standard populations (see Methods)

Table A16 – Cancer of the Small intestine, incl. duodenum (152)

South Australia Year – 2002	Incidence						Mortality					
	New cases			Rates			Deaths			Rates		
	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons
0–4	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
5–9	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
10–14	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
15–19	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
20–24	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
25–29	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
30–34	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
35–39	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
40–44	1	1	2	1.7	1.7	1.7	0	0	0	0.0	0.0	0.0
45–49	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
50–54	1	0	1	1.9	0.0	1.0	0	0	0	0.0	0.0	0.0
55–59	0	2	2	0.0	4.5	2.3	0	1	1	0.0	2.3	1.1
60–64	4	1	5	11.7	2.9	7.3	1	2	3	2.9	5.8	4.4
65–69	2	2	4	6.9	6.6	6.8	1	0	1	3.5	0.0	1.7
70–74	1	3	4	3.8	10.1	7.1	0	3	3	0.0	10.1	5.3
75–79	4	5	9	18.5	18.0	18.2	1	0	1	4.6	0.0	2.0
80–84	0	2	2	0.0	9.9	6.1	2	1	3	15.6	5.0	9.1
85+	1	2	3	12.5	10.9	11.4	0	1	1	0.0	5.5	3.8
Total/Crude rate	14	18	32	1.9	2.3	2.1	5	8	13	0.7	1.0	0.9
AS Rate (Aust)				1.8	1.8	1.8				0.6	0.9	0.7
Confidence Interval (95%)				0.8 – 2.7	1.0 – 2.7	1.2 – 2.5				0.1 – 1.2	0.3 – 1.4	0.3 – 1.1
AS Rate (World)				1.3	1.3	1.3				0.4	0.6	0.5
Confidence Interval (95%)				0.6 – 2.0	0.7 – 1.8	0.9 – 1.7				0.1 – 0.8	0.2 – 1.1	0.2 – 0.8
Risk				1 in 767	1 in 776	1 in 768				1 in 3,123	1 in 1,105	1 in 1,599
PYLL										20	50	70
% of all cancers	0.3	0.5	0.4				0.3	0.5	0.4	0.3	0.5	0.4
Trend 1998–2002 (per annum)				5.4%	53.9%	20.4%				-11.1%	20.1%	-2.3%

Table A17 – Cancer of the Large intestine, excluding rectum (153)

South Australia Year – 2002	Incidence						Mortality					
	New cases			Rates			Deaths			Rates		
	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons
0–4	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
5–9	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
10–14	1	1	2	1.9	2.0	2.0	0	0	0	0.0	0.0	0.0
15–19	0	2	2	0.0	4.0	1.9	0	0	0	0.0	0.0	0.0
20–24	1	0	1	2.0	0.0	1.0	0	0	0	0.0	0.0	0.0
25–29	2	1	3	4.0	2.1	3.1	1	0	1	2.0	0.0	1.0
30–34	1	0	1	1.8	0.0	0.9	1	0	1	1.8	0.0	0.9
35–39	2	5	7	3.6	9.0	6.3	0	0	0	0.0	0.0	0.0
40–44	4	8	12	6.9	13.6	10.2	0	0	0	0.0	0.0	0.0
45–49	8	7	15	14.9	12.8	13.8	3	6	9	5.6	11.0	8.3
50–54	20	17	37	38.8	32.2	35.4	6	1	7	11.6	1.9	6.7
55–59	34	31	65	77.1	69.8	73.4	11	6	17	24.9	13.5	19.2
60–64	50	23	73	146.7	66.3	106.1	10	9	19	29.3	25.9	27.6
65–69	55	38	93	190.9	125.5	157.3	17	8	25	59.0	26.4	42.3
70–74	73	56	129	274.9	188.4	229.2	21	17	38	79.1	57.2	67.5
75–79	62	63	125	287.2	226.3	252.9	27	25	52	125.1	89.8	105.2
80–84	32	58	90	249.0	288.2	272.9	18	23	41	140.1	114.3	124.3
85+	28	71	99	349.0	388.4	376.4	23	38	61	286.7	207.9	231.9
Total/Crude rate	373	381	754	49.6	49.6	49.6	138	133	271	18.4	17.3	17.8
AS Rate (Aust)				47.8	38.7	43.2				18.3	12.7	15.1
Confidence Interval (95%)				42.9 – 52.7	34.7 – 42.6	40.1 – 46.3				15.2 – 21.4	10.5 – 14.9	13.3 – 17.0
AS Rate (World)				33.9	26.6	30.2				11.9	8.0	9.8
Confidence Interval (95%)				30.6 – 37.1	23.9 – 29.3	28.1 – 32.3				10.1 – 13.7	6.7 – 9.4	8.7 – 10.9
Risk				1 in 27	1 in 39	1 in 32				1 in 94	1 in 148	1 in 116
PYLL										805	508	1,313
% of all cancers	9.1	10.5	9.8				9.1	10.5	9.8	7.6	8.8	8.2
Trend 1998–2002 (per annum)				1.5%	1.7%	1.9%				3.0%	-1.7%	0.6%

Table A18 – Cancer of the Hepatic flexure (153.0)

South Australia Year – 2002	Incidence						Mortality					
	New cases			Rates			Deaths			Rates		
	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons
0–4	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
5–9	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
10–14	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
15–19	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
20–24	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
25–29	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
30–34	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
35–39	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
40–44	1	0	1	1.7	0.0	0.9	0	0	0	0.0	0.0	0.0
45–49	0	0	0	0.0	0.0	0.0	0	1	1	0.0	1.8	0.9
50–54	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
55–59	0	2	2	0.0	4.5	2.3	0	1	1	0.0	2.3	1.1
60–64	1	1	2	2.9	2.9	2.9	0	1	1	0.0	2.9	1.5
65–69	2	1	3	6.9	3.3	5.1	0	0	0	0.0	0.0	0.0
70–74	4	8	12	15.1	26.9	21.3	2	2	4	7.5	6.7	7.1
75–79	3	2	5	13.9	7.2	10.1	1	1	2	4.6	3.6	4.0
80–84	3	1	4	23.3	5.0	12.1	1	0	1	7.8	0.0	3.0
85+	1	4	5	12.5	21.9	19.0	0	2	2	0.0	10.9	7.6
Total/Crude rate	15	19	34	2.0	2.5	2.2	4	8	12	0.5	1.0	0.8
AS Rate (Aust)				1.9	1.9	1.9				0.5	0.8	0.7
Confidence Interval (95%)				1.0 – 2.9	1.1 – 2.8	1.3 – 2.6				0.0 – 1.0	0.2 – 1.4	0.3 – 1.1
AS Rate (World)				1.3	1.3	1.3				0.3	0.6	0.5
Confidence Interval (95%)				0.7 – 1.8	0.7 – 1.8	0.9 – 1.7				0.1 – 0.5	0.2 – 1.0	0.2 – 0.7
Risk				1 in 751	1 in 532	1 in 617				1 in 2,656	1 in 1,462	1 in 1,885
PYLL										5	63	68
% of all cancers	0.4	0.5	0.4				0.4	0.5	0.4	0.2	0.5	0.4
Trend 1998–2002 (per annum)				6.4%	14.4%	9.2%				-6.7%	1.4%	-3.8%

Note: All rates are expressed per 100,000 population. Age-standardised rates (AS Rate) use the Australian 2001 and New World Standard populations (see Methods)

Table A19 – Cancer of the Transverse colon (153.1)

South Australia Year – 2002	Incidence						Mortality					
	New cases			Rates			Deaths			Rates		
	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons
0–4	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
5–9	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
10–14	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
15–19	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
20–24	1	0	1	2.0	0.0	1.0	0	0	0	0.0	0.0	0.0
25–29	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
30–34	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
35–39	1	0	1	1.8	0.0	0.9	0	0	0	0.0	0.0	0.0
40–44	1	0	1	1.7	0.0	0.9	0	0	0	0.0	0.0	0.0
45–49	1	0	1	1.9	0.0	0.9	0	0	0	0.0	0.0	0.0
50–54	4	2	6	7.8	3.8	5.7	1	0	1	1.9	0.0	1.0
55–59	1	2	3	2.3	4.5	3.4	0	0	0	0.0	0.0	0.0
60–64	6	4	10	17.6	11.5	14.5	0	3	3	0.0	8.6	4.4
65–69	2	6	8	6.9	19.8	13.5	3	1	4	10.4	3.3	6.8
70–74	5	2	7	18.8	6.7	12.4	2	2	4	7.5	6.7	7.1
75–79	7	7	14	32.4	25.1	28.3	2	3	5	9.3	10.8	10.1
80–84	3	2	5	23.3	9.9	15.2	0	2	2	0.0	9.9	6.1
85+	3	6	9	37.4	32.8	34.2	3	2	5	37.4	10.9	19.0
Total/Crude rate	35	31	66	4.7	4.0	4.3	11	13	24	1.5	1.7	1.6
AS Rate (Aust)				4.6	3.2	3.8				1.5	1.3	1.4
Confidence Interval (95%)				3.0 – 6.1	2.0 – 4.3	2.9 – 4.7				0.6 – 2.4	0.6 – 2.0	0.8 – 1.9
AS Rate (World)				3.2	2.2	2.7				0.9	0.9	0.9
Confidence Interval (95%)				2.2 – 4.3	1.4 – 3.1	2.1 – 3.4				0.4 – 1.5	0.4 – 1.4	0.5 – 1.2
Risk				1 in 330	1 in 432	1 in 375				1 in 1,007	1 in 1,072	1 in 1,043
PYLL										50	50	100
% of all cancers	0.9	0.9	0.9				0.9	0.9	0.9	0.6	0.9	0.7
Trend 1998–2002 (per annum)				7.4%	-2.5%	1.9%				3.5%	4.4%	3.6%

Table A20 – Cancer of the Descending colon (153.2)

South Australia Year – 2002	Incidence						Mortality					
	New cases			Rates			Deaths			Rates		
	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons
0–4	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
5–9	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
10–14	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
15–19	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
20–24	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
25–29	0	1	1	0.0	2.1	1.0	0	0	0	0.0	0.0	0.0
30–34	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
35–39	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
40–44	0	1	1	0.0	1.7	0.9	0	0	0	0.0	0.0	0.0
45–49	0	1	1	0.0	1.8	0.9	0	1	1	0.0	1.8	0.9
50–54	1	1	2	1.9	1.9	1.9	0	0	0	0.0	0.0	0.0
55–59	3	2	5	6.8	4.5	5.6	0	0	0	0.0	0.0	0.0
60–64	3	1	4	8.8	2.9	5.8	0	1	1	0.0	2.9	1.5
65–69	3	0	3	10.4	0.0	5.1	1	1	2	3.5	3.3	3.4
70–74	5	3	8	18.8	10.1	14.2	2	1	3	7.5	3.4	5.3
75–79	8	3	11	37.1	10.8	22.3	2	1	3	9.3	3.6	6.1
80–84	3	1	4	23.3	5.0	12.1	1	1	2	7.8	5.0	6.1
85+	2	4	6	24.9	21.9	22.8	0	0	0	0.0	0.0	0.0
Total/Crude rate	28	18	46	3.7	2.3	3.0	6	6	12	0.8	0.8	0.8
AS Rate (Aust)				3.6	1.9	2.6				0.7	0.7	0.7
Confidence Interval (95%)				2.2 – 4.9	1.0 – 2.8	1.9 – 3.4				0.1 – 1.4	0.1 – 1.2	0.3 – 1.1
AS Rate (World)				2.4	1.4	1.8				0.5	0.5	0.5
Confidence Interval (95%)				1.6 – 3.2	0.7 – 2.0	1.3 – 2.3				0.2 – 0.8	0.1 – 0.9	0.2 – 0.7
Risk				1 in 428	1 in 800	1 in 564				1 in 1,818	1 in 1,759	1 in 1,804
PYLL										13	50	63
% of all cancers	0.7	0.5	0.6				0.7	0.5	0.6	0.3	0.4	0.4
Trend 1998–2002 (per annum)				6.4%	5.6%	7.0%				-7.4%	10.9%	-1.3%

Table A21 – Cancer of the Sigmoid colon (153.3)

South Australia Year – 2002	Incidence						Mortality					
	New cases			Rates			Deaths			Rates		
	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons
0–4	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
5–9	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
10–14	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
15–19	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
20–24	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
25–29	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
30–34	1	0	1	1.8	0.0	0.9	0	0	0	0.0	0.0	0.0
35–39	0	3	3	0.0	5.4	2.7	0	0	0	0.0	0.0	0.0
40–44	0	4	4	0.0	6.8	3.4	0	0	0	0.0	0.0	0.0
45–49	5	4	9	9.3	7.3	8.3	1	2	3	1.9	3.7	2.8
50–54	9	8	17	17.4	15.2	16.3	2	0	2	3.9	0.0	1.9
55–59	13	16	29	29.5	36.0	32.8	4	1	5	9.1	2.3	5.6
60–64	19	3	22	55.7	8.6	32.0	2	0	2	5.9	0.0	2.9
65–69	17	9	26	59.0	29.7	44.0	5	3	8	17.4	9.9	13.5
70–74	21	18	39	79.1	60.6	69.3	4	6	10	15.1	20.2	17.8
75–79	16	14	30	74.1	50.3	60.7	9	6	15	41.7	21.6	30.3
80–84	2	9	11	15.6	44.7	33.4	5	1	6	38.9	5.0	18.2
85+	4	17	21	49.9	93.0	79.8	8	9	17	99.7	49.2	64.6
Total/Crude rate	107	105	212	14.2	13.7	14.0	40	28	68	5.3	3.6	4.5
AS Rate (Aust)				13.5	11.1	12.4				5.4	2.7	3.8
Confidence Interval (95%)				10.9 – 16.0	9.0 – 13.3	10.7 – 14.0				3.7 – 7.0	1.7 – 3.7	2.9 – 4.7
AS Rate (World)				10.1	8.0	9.1				3.4	1.7	2.4
Confidence Interval (95%)				8.3 – 12.0	6.5 – 9.5	7.9 – 10.3				2.4 – 4.3	1.1 – 2.4	1.9 – 3.0
Risk				1 in 80	1 in 118	1 in 96				1 in 377	1 in 556	1 in 450
PYLL										215	110	325
% of all cancers	2.6	2.9	2.8				2.6	2.9	2.8	2.2	1.9	2.0
Trend 1998–2002 (per annum)				-5.7%	1.3%	-2.4%				3.0%	-5.2%	-0.2%

Note: All rates are expressed per 100,000 population. Age-standardised rates (AS Rate) use the Australian 2001 and New World Standard populations (see Methods)

Table A22 – Cancer of the Caecum (153.4)

South Australia Year – 2002	Incidence						Mortality					
	New cases			Rates			Deaths			Rates		
	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons
0–4	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
5–9	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
10–14	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
15–19	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
20–24	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
25–29	1	0	1	2.0	0.0	1.0	1	0	1	2.0	0.0	1.0
30–34	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
35–39	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
40–44	0	1	1	0.0	1.7	0.9	0	0	0	0.0	0.0	0.0
45–49	1	1	2	1.9	1.8	1.8	1	0	1	1.9	0.0	0.9
50–54	3	4	7	5.8	7.6	6.7	0	0	0	0.0	0.0	0.0
55–59	11	4	15	24.9	9.0	16.9	3	2	5	6.8	4.5	5.6
60–64	13	5	18	38.1	14.4	26.2	5	0	5	14.7	0.0	7.3
65–69	11	8	19	38.2	26.4	32.1	3	0	3	10.4	0.0	5.1
70–74	17	14	31	64.0	47.1	55.1	5	3	8	18.8	10.1	14.2
75–79	16	19	35	74.1	68.2	70.8	7	8	15	32.4	28.7	30.3
80–84	10	25	35	77.8	124.2	106.1	3	9	12	23.3	44.7	36.4
85+	9	20	29	112.2	109.4	110.3	4	7	11	49.9	38.3	41.8
Total/Crude rate	92	101	193	12.2	13.2	12.7	32	29	61	4.3	3.8	4.0
AS Rate (Aust)				11.9	9.8	10.9				4.2	2.6	3.4
Confidence Interval (95%)				9.4 – 14.3	7.8 – 11.7	9.3 – 12.4				2.7 – 5.6	1.7 – 3.6	2.5 – 4.2
AS Rate (World)				8.2	6.2	7.2				2.8	1.5	2.2
Confidence Interval (95%)				6.6 – 9.7	5.0 – 7.4	6.2 – 8.2				1.9 – 3.8	1.1 – 1.9	1.7 – 2.7
Risk				1 in 115	1 in 186	1 in 143				1 in 367	1 in 1,371	1 in 586
PYLL										225	43	268
% of all cancers	2.2	2.8	2.5				2.2	2.8	2.5	1.8	1.9	1.8
Trend 1998–2002 (per annum)				5.2%	-2.3%	1.3%				1.7%	-6.7%	-2.8%

Table A23 – Cancer of the Appendix (153.5)

South Australia Year – 2002	Incidence						Mortality					
	New cases			Rates			Deaths			Rates		
	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons
0–4	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
5–9	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
10–14	1	0	1	1.9	0.0	1.0	0	0	0	0.0	0.0	0.0
15–19	0	2	2	0.0	4.0	1.9	0	0	0	0.0	0.0	0.0
20–24	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
25–29	1	0	1	2.0	0.0	1.0	0	0	0	0.0	0.0	0.0
30–34	0	0	0	0.0	0.0	0.0	1	0	1	1.8	0.0	0.9
35–39	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
40–44	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
45–49	0	0	0	0.0	0.0	0.0	0	1	1	0.0	1.8	0.9
50–54	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
55–59	1	1	2	2.3	2.3	2.3	1	0	1	2.3	0.0	1.1
60–64	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
65–69	0	2	2	0.0	6.6	3.4	0	0	0	0.0	0.0	0.0
70–74	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
75–79	0	0	0	0.0	0.0	0.0	0	1	1	0.0	3.6	2.0
80–84	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
85+	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
Total/Crude rate	3	5	8	0.4	0.7	0.5	2	2	4	0.3	0.3	0.3
AS Rate (Aust)				0.4	0.6	0.5				0.3	0.2	0.2
Confidence Interval (95%)				0.0 – 0.9	0.1 – 1.2	0.2 – 0.9				0.0 – 0.6	0.0 – 0.5	0.0 – 0.5
AS Rate (World)				0.4	0.6	0.5				0.2	0.2	0.2
Confidence Interval (95%)				0.0 – 0.9	0.0 – 1.2	0.1 – 0.9				0.0 – 0.5	0.0 – 0.4	0.0 – 0.4
Risk				1 in 3,217	1 in 1,561	1 in 2,084				1 in 4,901	1 in 10,944	1 in 6,737
PYLL										60	28	88
% of all cancers	0.1	0.1	0.1				0.1	0.1	0.1	0.1	0.1	0.1
Trend 1998–2002 (per annum)				7.7%	5.4%	5.7%				240.0%	46.0%	101.4%

Table A24 – Cancer of the Ascending colon (153.6)

South Australia Year – 2002	Incidence						Mortality					
	New cases			Rates			Deaths			Rates		
	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons
0–4	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
5–9	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
10–14	0	1	1	0.0	2.0	1.0	0	0	0	0.0	0.0	0.0
15–19	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
20–24	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
25–29	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
30–34	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
35–39	0	2	2	0.0	3.6	1.8	0	0	0	0.0	0.0	0.0
40–44	1	0	1	1.7	0.0	0.9	0	0	0	0.0	0.0	0.0
45–49	1	0	1	1.9	0.0	0.9	0	1	1	0.0	1.8	0.9
50–54	2	0	2	3.9	0.0	1.9	2	1	3	3.9	1.9	2.9
55–59	4	2	6	9.1	4.5	6.8	1	2	3	2.3	4.5	3.4
60–64	5	7	12	14.7	20.2	17.4	1	3	4	2.9	8.6	5.8
65–69	15	8	23	52.1	26.4	38.9	3	1	4	10.4	3.3	6.8
70–74	14	10	24	52.7	33.6	42.6	3	2	5	11.3	6.7	8.9
75–79	10	13	23	46.3	46.7	46.5	2	3	5	9.3	10.8	10.1
80–84	8	14	22	62.3	69.6	66.7	6	6	12	46.7	29.8	36.4
85+	0	8	8	0.0	43.8	30.4	4	7	11	49.9	38.3	41.8
Total/Crude rate	60	65	125	8.0	8.5	8.2	22	26	48	2.9	3.4	3.2
AS Rate (Aust)				7.5	6.6	7.1				3.0	2.5	2.7
Confidence Interval (95%)				5.6 – 9.4	5.0 – 8.2	5.9 – 8.4				1.7 – 4.2	1.5 – 3.5	1.9 – 3.4
AS Rate (World)				5.4	4.5	5.0				1.8	1.6	1.7
Confidence Interval (95%)				4.1 – 6.6	3.4 – 5.6	4.1 – 5.8				1.2 – 2.5	1.0 – 2.3	1.3 – 2.2
Risk				1 in 148	1 in 222	1 in 179				1 in 650	1 in 744	1 in 699
PYLL										105	135	240
% of all cancers	1.5	1.8	1.6				1.5	1.8	1.6	1.2	1.7	1.4
Trend 1998–2002 (per annum)				3.1%	2.5%	3.7%				10.0%	0.7%	3.4%

Note: All rates are expressed per 100,000 population. Age-standardised rates (AS Rate) use the Australian 2001 and New World Standard populations (see Methods)

Table A25 – Cancer of the Splenic flexure (153.7)

South Australia Year – 2002	Incidence						Mortality					
	New cases			Rates			Deaths			Rates		
	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons
0–4	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
5–9	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
10–14	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
15–19	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
20–24	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
25–29	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
30–34	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
35–39	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
40–44	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
45–49	0	1	1	0.0	1.8	0.9	1	0	1	1.9	0.0	0.9
50–54	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
55–59	1	2	3	2.3	4.5	3.4	0	0	0	0.0	0.0	0.0
60–64	0	1	1	0.0	2.9	1.5	1	1	2	2.9	2.9	2.9
65–69	2	2	4	6.9	6.6	6.8	2	1	3	6.9	3.3	5.1
70–74	3	1	4	11.3	3.4	7.1	0	1	1	0.0	3.4	1.8
75–79	2	3	5	9.3	10.8	10.1	2	0	2	9.3	0.0	4.0
80–84	1	2	3	7.8	9.9	9.1	0	0	0	0.0	0.0	0.0
85+	2	5	7	24.9	27.4	26.6	0	3	3	0.0	16.4	11.4
Total/Crude rate	11	17	28	1.5	2.2	1.8	6	6	12	0.8	0.8	0.8
AS Rate (Aust)				1.5	1.7	1.6				0.7	0.6	0.7
Confidence Interval (95%)				0.6 – 2.3	0.9 – 2.5	1.0 – 2.1				0.1 – 1.3	0.1 – 1.0	0.3 – 1.1
AS Rate (World)				0.9	1.1	1.0				0.6	0.4	0.5
Confidence Interval (95%)				0.4 – 1.4	0.6 – 1.6	0.7 – 1.4				0.1 – 1.0	0.0 – 0.7	0.2 – 0.8
Risk				1 in 976	1 in 1,043	1 in 1,019				1 in 1,704	1 in 2,095	1 in 1,873
PYLL										55	23	78
% of all cancers	0.3	0.5	0.4				0.3	0.5	0.4	0.3	0.4	0.4
Trend 1998–2002 (per annum)				-1.5%	16.9%	6.6%				-3.6%	-1.9%	-2.7%

Table A26 – Cancer of other specified sites of the Colon (153.8)

South Australia Year – 2002	Incidence						Mortality					
	New cases			Rates			Deaths			Rates		
	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons
0–4	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
5–9	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
10–14	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
15–19	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
20–24	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
25–29	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
30–34	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
35–39	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
40–44	1	1	2	1.7	1.7	1.7	0	0	0	0.0	0.0	0.0
45–49	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
50–54	1	1	2	1.9	1.9	1.9	0	0	0	0.0	0.0	0.0
55–59	0	0	0	0.0	0.0	0.0	1	0	1	2.3	0.0	1.1
60–64	1	1	2	2.9	2.9	2.9	0	0	0	0.0	0.0	0.0
65–69	2	1	3	6.9	3.3	5.1	0	0	0	0.0	0.0	0.0
70–74	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
75–79	0	1	1	0.0	3.6	2.0	1	1	2	4.6	3.6	4.0
80–84	0	1	1	0.0	5.0	3.0	0	0	0	0.0	0.0	0.0
85+	3	0	3	37.4	0.0	11.4	0	1	1	0.0	5.5	3.8
Total/Crude rate	8	6	14	1.1	0.8	0.9	2	2	4	0.3	0.3	0.3
AS Rate (Aust)				1.1	0.7	0.8				0.2	0.2	0.2
Confidence Interval (95%)				0.3 – 1.9	0.1 – 1.2	0.4 – 1.3				0.0 – 0.6	0.0 – 0.4	0.0 – 0.4
AS Rate (World)				0.8	0.5	0.6				0.2	0.1	0.1
Confidence Interval (95%)				0.3 – 1.3	0.1 – 0.9	0.3 – 0.9				0.0 – 0.4	0.0 – 0.2	0.0 – 0.2
Risk				1 in 1,479	1 in 2,046	1 in 1,724				1 in 8,821	N/A	1 in 17,705
PYLL										18	0	18
% of all cancers	0.2	0.2	0.2				0.2	0.2	0.2	0.1	0.1	0.1
Trend 1998–2002 (per annum)				50.8%	28.6%	31.4%				6.2%	18.1%	8.8%

Table A27 – Cancer of the Colon, unspecified (153.9)

South Australia Year – 2002	Incidence						Mortality					
	New cases			Rates			Deaths			Rates		
	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons
0–4	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
5–9	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
10–14	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
15–19	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
20–24	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
25–29	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
30–34	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
35–39	1	0	1	1.8	0.0	0.9	0	0	0	0.0	0.0	0.0
40–44	0	1	1	0.0	1.7	0.9	0	0	0	0.0	0.0	0.0
45–49	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
50–54	0	1	1	0.0	1.9	1.0	1	0	1	1.9	0.0	1.0
55–59	0	0	0	0.0	0.0	0.0	1	0	1	2.3	0.0	1.1
60–64	2	0	2	5.9	0.0	2.9	1	0	1	2.9	0.0	1.5
65–69	1	1	2	3.5	3.3	3.4	0	1	1	0.0	3.3	1.7
70–74	4	0	4	15.1	0.0	7.1	3	0	3	11.3	0.0	5.3
75–79	0	1	1	0.0	3.6	2.0	1	1	2	4.6	3.6	4.0
80–84	2	3	5	15.6	14.9	15.2	2	4	6	15.6	19.9	18.2
85+	4	7	11	49.9	38.3	41.8	4	7	11	49.9	38.3	41.8
Total/Crude rate	14	14	28	1.9	1.8	1.8	13	13	26	1.7	1.7	1.7
AS Rate (Aust)				1.9	1.2	1.6				1.8	1.1	1.4
Confidence Interval (95%)				0.9 – 3.0	0.6 – 1.9	1.0 – 2.1				0.8 – 2.8	0.5 – 1.7	0.9 – 2.0
AS Rate (World)				1.2	0.7	1.0				1.1	0.6	0.8
Confidence Interval (95%)				0.6 – 1.8	0.4 – 1.1	0.6 – 1.3				0.6 – 1.6	0.3 – 0.8	0.5 – 1.1
Risk				1 in 764	1 in 2,901	1 in 1,242				1 in 1,085	1 in 6,058	1 in 1,894
PYLL										60	8	68
% of all cancers	0.3	0.4	0.4				0.3	0.4	0.4	0.7	0.9	0.8
Trend 1998–2002 (per annum)				27.1%	12.4%	17.4%				19.4%	9.7%	14.6%

Note: All rates are expressed per 100,000 population. Age-standardised rates (AS Rate) use the Australian 2001 and New World Standard populations (see Methods)

Table A28 – Cancer of the Rectum, rectosigmoid junction and anal canal (154)

South Australia Year – 2002	Incidence						Mortality					
	New cases			Rates			Deaths			Rates		
	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons
0-4	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
5-9	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
10-14	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
15-19	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
20-24	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
25-29	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
30-34	2	0	2	3.6	0.0	1.8	1	0	1	1.8	0.0	0.9
35-39	4	1	5	7.2	1.8	4.5	0	2	2	0.0	3.6	1.8
40-44	3	3	6	5.1	5.1	5.1	0	1	1	0.0	1.7	0.9
45-49	6	7	13	11.2	12.8	12.0	3	1	4	5.6	1.8	3.7
50-54	25	12	37	48.4	22.7	35.4	9	4	13	17.4	7.6	12.5
55-59	19	19	38	43.1	42.8	42.9	6	7	13	13.6	15.8	14.7
60-64	32	14	46	93.9	40.3	66.9	13	3	16	38.1	8.6	23.3
65-69	39	17	56	135.3	56.1	94.7	14	3	17	48.6	9.9	28.8
70-74	56	30	86	210.9	100.9	152.8	11	9	20	41.4	30.3	35.5
75-79	36	31	67	166.8	111.3	135.6	11	8	19	51.0	28.7	38.4
80-84	23	17	40	179.0	84.5	121.3	13	12	25	101.2	59.6	75.8
85+	9	22	31	112.2	120.4	117.9	6	21	27	74.8	114.9	102.7
Total/Crude rate	254	173	427	33.8	22.5	28.1	87	71	158	11.6	9.3	10.4
AS Rate (Aust)				32.2	18.2	24.8				11.2	6.9	9.0
Confidence Interval (95%)				28.2 – 36.2	15.5 – 21.0	22.4 – 27.2				8.8 – 13.6	5.3 – 8.6	7.6 – 10.4
AS Rate (World)				23.4	13.0	17.9				7.9	4.6	6.2
Confidence Interval (95%)				20.7 – 26.1	11.1 – 14.9	16.3 – 19.6				6.4 – 9.5	3.5 – 5.6	5.3 – 7.1
Risk				1 in 36	1 in 71	1 in 49				1 in 121	1 in 253	1 in 164
PYLL										728	430	1,158
% of all cancers	6.2	4.8	5.5				6.2	4.8	5.5	4.8	4.7	4.8
Trend 1998-2002 (per annum)				-0.2%	0.4%	0.1%				-4.7%	-4.3%	-4.2%

Table A29 – Cancer of the Rectosigmoid junction (154.0)

South Australia Year – 2002	Incidence						Mortality					
	New cases			Rates			Deaths			Rates		
	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons
0-4	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
5-9	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
10-14	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
15-19	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
20-24	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
25-29	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
30-34	0	0	0	0.0	0.0	0.0	1	0	1	1.8	0.0	0.9
35-39	3	0	3	5.4	0.0	2.7	0	0	0	0.0	0.0	0.0
40-44	1	1	2	1.7	1.7	1.7	0	0	0	0.0	0.0	0.0
45-49	0	3	3	0.0	5.5	2.8	1	0	1	1.9	0.0	0.9
50-54	2	4	6	3.9	7.6	5.7	2	1	3	3.9	1.9	2.9
55-59	3	4	7	6.8	9.0	7.9	1	2	3	2.3	4.5	3.4
60-64	8	2	10	23.5	5.8	14.5	5	1	6	14.7	2.9	8.7
65-69	12	3	15	41.6	9.9	25.4	3	0	3	10.4	0.0	5.1
70-74	11	9	20	41.4	30.3	35.5	3	5	8	11.3	16.8	14.2
75-79	13	7	20	60.2	25.1	40.5	4	4	8	18.5	14.4	16.2
80-84	6	11	17	46.7	54.7	51.6	5	3	8	38.9	14.9	24.3
85+	4	9	13	49.9	49.2	49.4	0	5	5	0.0	27.4	19.0
Total/Crude rate	63	53	116	8.4	6.9	7.6	25	21	46	3.3	2.7	3.0
AS Rate (Aust)				8.1	5.4	6.6				3.2	2.0	2.6
Confidence Interval (95%)				6.1 – 10.1	3.9 – 6.8	5.4 – 7.8				1.9 – 4.4	1.2 – 2.9	1.9 – 3.4
AS Rate (World)				5.7	3.6	4.6				2.3	1.3	1.8
Confidence Interval (95%)				4.4 – 7.0	2.7 – 4.6	3.8 – 5.4				1.4 – 3.1	0.8 – 1.8	1.3 – 2.3
Risk				1 in 161	1 in 287	1 in 208				1 in 433	1 in 767	1 in 554
PYLL										225	83	308
% of all cancers	1.5	1.5	1.5				1.5	1.5	1.5	1.4	1.4	1.4
Trend 1998-2002 (per annum)				-6.2%	-0.5%	-3.7%				-2.4%	-7.0%	-3.6%

Table A30 – Cancer of the Rectum (154.1)

South Australia Year – 2002	Incidence						Mortality					
	New cases			Rates			Deaths			Rates		
	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons
0-4	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
5-9	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
10-14	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
15-19	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
20-24	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
25-29	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
30-34	2	0	2	3.6	0.0	1.8	0	0	0	0.0	0.0	0.0
35-39	1	1	2	1.8	1.8	1.8	0	2	2	0.0	3.6	1.8
40-44	2	2	4	3.4	3.4	3.4	0	1	1	0.0	1.7	0.9
45-49	6	4	10	11.2	7.3	9.2	2	1	3	3.7	1.8	2.8
50-54	23	8	31	44.6	15.2	29.7	7	3	10	13.6	5.7	9.6
55-59	16	13	29	36.3	29.3	32.8	5	4	9	11.3	9.0	10.2
60-64	22	11	33	64.6	31.7	48.0	8	2	10	23.5	5.8	14.5
65-69	27	12	39	93.7	39.6	66.0	11	3	14	38.2	9.9	23.7
70-74	45	21	66	169.4	70.6	117.3	8	4	12	30.1	13.5	21.3
75-79	23	22	45	106.6	79.0	91.0	7	4	11	32.4	14.4	22.3
80-84	16	6	22	124.5	29.8	66.7	8	8	16	62.3	39.8	48.5
85+	5	13	18	62.3	71.1	68.4	6	16	22	74.8	87.5	83.6
Total/Crude rate	188	113	301	25.0	14.7	19.8	62	48	110	8.3	6.3	7.2
AS Rate (Aust)				23.7	12.1	17.6				8.0	4.7	6.3
Confidence Interval (95%)				20.3 – 27.1	9.8 – 14.4	15.6 – 19.6				6.0 – 10.0	3.3 – 6.0	5.1 – 7.5
AS Rate (World)				17.4	8.8	12.9				5.6	3.1	4.3
Confidence Interval (95%)				15.1 – 19.7	7.2 – 10.3	11.5 – 14.3				4.3 – 7.0	2.2 – 4.0	3.5 – 5.1
Risk				1 in 47	1 in 101	1 in 65				1 in 167	1 in 393	1 in 237
PYLL										503	330	833
% of all cancers	4.6	3.1	3.9				4.6	3.1	3.9	3.4	3.2	3.3
Trend 1998-2002 (per annum)				3.1%	1.4%	2.3%				-4.9%	-3.1%	-4.1%

Note: All rates are expressed per 100,000 population. Age-standardised rates (AS Rate) use the Australian 2001 and New World Standard populations (see Methods)

Table A31 – Cancer of the Anal canal (154.2)

South Australia Year – 2002	Incidence						Mortality					
	New cases			Rates			Deaths			Rates		
	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons
0-4	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
5-9	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
10-14	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
15-19	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
20-24	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
25-29	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
30-34	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
35-39	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
40-44	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
45-49	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
50-54	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
55-59	0	1	1	0.0	2.3	1.1	0	1	1	0.0	2.3	1.1
60-64	1	0	1	2.9	0.0	1.5	0	0	0	0.0	0.0	0.0
65-69	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
70-74	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
75-79	0	1	1	0.0	3.6	2.0	0	0	0	0.0	0.0	0.0
80-84	0	0	0	0.0	0.0	0.0	0	1	1	0.0	5.0	3.0
85+	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
Total/Crude rate	1	2	3	0.1	0.3	0.2	0	2	2	0.0	0.3	0.1
AS Rate (Aust)				0.1	0.2	0.2				0.0	0.2	0.1
Confidence Interval (95%)				0.0 – 0.4	0.0 – 0.5	0.0 – 0.4				0.0 – 0.0	0.0 – 0.5	0.0 – 0.3
AS Rate (World)				0.1	0.2	0.1				0.0	0.1	0.1
Confidence Interval (95%)				0.0 – 0.3	0.0 – 0.3	0.0 – 0.3				0.0 – 0.0	0.0 – 0.3	0.0 – 0.2
Risk				1 in 6,817	1 in 8,885	1 in 7,743				N/A	1 in 8,885	1 in 17,705
PYLL										0	18	18
% of all cancers	0.0	0.1	0.0				0.0	0.1	0.0	0.0	0.1	0.1
Trend 1998–2002 (per annum)				-16.1%	-7.5%	-9.3%				-33.3%	2.8%	-16.8%

Table A32 – Cancer of the Anorectal junction (154.8)

South Australia Year – 2002	Incidence						Mortality					
	New cases			Rates			Deaths			Rates		
	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons
0-4	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
5-9	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
10-14	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
15-19	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
20-24	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
25-29	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
30-34	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
35-39	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
40-44	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
45-49	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
50-54	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
55-59	0	1	1	0.0	2.3	1.1	0	0	0	0.0	0.0	0.0
60-64	1	1	2	2.9	2.9	2.9	0	0	0	0.0	0.0	0.0
65-69	0	2	2	0.0	6.6	3.4	0	0	0	0.0	0.0	0.0
70-74	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
75-79	0	1	1	0.0	3.6	2.0	0	0	0	0.0	0.0	0.0
80-84	1	0	1	7.8	0.0	3.0	0	0	0	0.0	0.0	0.0
85+	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
Total/Crude rate	2	5	7	0.3	0.7	0.5	0	0	0	0.0	0.0	0.0
AS Rate (Aust)				0.3	0.6	0.4				0.0	0.0	0.0
Confidence Interval (95%)				0.0 – 0.6	0.1 – 1.1	0.1 – 0.7				0.0 – 0.0	0.0 – 0.0	0.0 – 0.0
AS Rate (World)				0.2	0.5	0.3				0.0	0.0	0.0
Confidence Interval (95%)				0.0 – 0.4	0.1 – 0.9	0.1 – 0.6				0.0 – 0.0	0.0 – 0.0	0.0 – 0.0
Risk				1 in 6,817	1 in 1,705	1 in 2,696				N/A	N/A	N/A
PYLL										0	0	0
% of all cancers	0.0	0.1	0.1				0.0	0.1	0.1	0.0	0.0	0.0
Trend 1998–2002 (per annum)				-2.0%	-1.5%	-2.7%				-2.1%	-6.0%	-5.6%

Table A33 – Cancer of the Colon and Rectum (including anus) (153-154)

South Australia Year – 2002	Incidence						Mortality					
	New cases			Rates			Deaths			Rates		
	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons
0-4	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
5-9	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
10-14	1	1	2	1.9	2.0	2.0	0	0	0	0.0	0.0	0.0
15-19	0	2	2	0.0	4.0	1.9	0	0	0	0.0	0.0	0.0
20-24	1	0	1	2.0	0.0	1.0	0	0	0	0.0	0.0	0.0
25-29	2	1	3	4.0	2.1	3.1	1	0	1	2.0	0.0	1.0
30-34	3	0	3	5.4	0.0	2.7	2	0	2	3.6	0.0	1.8
35-39	6	6	12	10.7	10.8	10.8	0	2	2	0.0	3.6	1.8
40-44	7	11	18	12.0	18.7	15.4	0	1	1	0.0	1.7	0.9
45-49	14	14	28	26.1	25.6	25.8	6	7	13	11.2	12.8	12.0
50-54	45	29	74	87.2	54.9	70.9	15	5	20	29.1	9.5	19.2
55-59	53	50	103	120.2	112.6	116.4	17	13	30	38.5	29.3	33.9
60-64	82	37	119	240.6	106.6	173.0	23	12	35	67.5	34.6	50.9
65-69	94	55	149	326.2	181.6	252.1	31	11	42	107.6	36.3	71.1
70-74	129	86	215	485.7	289.3	382.0	32	26	58	120.5	87.5	103.1
75-79	98	94	192	454.0	337.6	388.5	38	33	71	176.1	118.5	143.6
80-84	55	75	130	428.0	372.7	394.2	31	35	66	241.2	173.9	200.2
85+	37	93	130	461.2	508.8	494.3	29	59	88	361.5	322.8	334.6
Total/Crude rate	627	547	1,174	83.5	71.3	77.8	225	202	427	29.9	26.3	28.2
AS Rate (Aust)				80.0	56.9	68.0				29.5	19.6	24.2
Confidence Interval (95%)				73.7 – 86.3	52.1 – 61.7	64.1 – 71.9				25.6 – 33.4	16.9 – 22.4	21.9 – 26.5
AS Rate (World)				57.2	39.6	48.1				19.8	12.6	16.0
Confidence Interval (95%)				53.0 – 61.5	36.3 – 42.9	45.5 – 50.8				17.4 – 22.2	10.9 – 14.3	14.5 – 17.4
Risk				1 in 16	1 in 25	1 in 19				1 in 53	1 in 93	1 in 68
PYLL										1,533	938	2,470
% of all cancers	15.3	15.1	15.3				15.3	15.1	15.3	12.4	13.4	12.9
Trend 1998–2002 (per annum)				0.8%	1.3%	1.2%				-0.4%	-2.6%	-1.3%

Note: All rates are expressed per 100,000 population. Age-standardised rates (AS Rate) use the Australian 2001 and New World Standard populations (see Methods)

Table A34 – Cancer of the Liver & intrahepatic bile ducts (primary) (155)

South Australia Year – 2002	Incidence						Mortality					
	New cases			Rates			Deaths			Rates		
	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons
0–4	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
5–9	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
10–14	0	0	0	0.0	0.0	0.0	0	1	1	0.0	2.0	1.0
15–19	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
20–24	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
25–29	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
30–34	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
35–39	2	0	2	3.6	0.0	1.8	0	0	0	0.0	0.0	0.0
40–44	1	0	1	1.7	0.0	0.9	0	0	0	0.0	0.0	0.0
45–49	4	0	4	7.5	0.0	3.7	1	1	2	1.9	1.8	1.8
50–54	6	0	6	11.6	0.0	5.7	3	1	4	5.8	1.9	3.8
55–59	3	1	4	6.8	2.3	4.5	2	1	3	4.5	2.3	3.4
60–64	4	2	6	11.7	5.8	8.7	1	0	1	2.9	0.0	1.5
65–69	5	2	7	17.4	6.6	11.8	3	2	5	10.4	6.6	8.5
70–74	8	3	11	30.1	10.1	19.5	9	2	11	33.9	6.7	19.5
75–79	6	6	12	27.8	21.6	24.3	3	5	8	13.9	18.0	16.2
80–84	6	1	7	46.7	5.0	21.2	5	3	8	38.9	14.9	24.3
85+	6	2	8	74.8	10.9	30.4	5	3	8	62.3	16.4	30.4
Total/Crude rate	51	17	68	6.8	2.2	4.5	32	19	51	4.3	2.5	3.4
AS Rate (Aust)				6.7	1.7	3.9				4.2	1.9	2.9
Confidence Interval (95%)				4.9 – 8.6	0.9 – 2.6	3.0 – 4.9				2.8 – 5.7	1.0 – 2.8	2.1 – 3.7
AS Rate (World)				4.7	1.2	2.8				2.7	1.3	1.9
Confidence Interval (95%)				3.5 – 5.8	0.6 – 1.7	2.1 – 3.4				1.9 – 3.6	0.7 – 2.0	1.4 – 2.4
Risk				1 in 222	1 in 810	1 in 353				1 in 337	1 in 937	1 in 507
PYLL										188	150	338
% of all cancers	1.2	0.5	0.9				1.2	0.5	0.9	1.8	1.3	1.5
Trend 1998–2002 (per annum)				13.4%	5.2%	10.4%				8.7%	7.6%	7.7%

Table A35 – Cancer of the Gallbladder & extrahepatic bile ducts (156)

South Australia Year – 2002	Incidence						Mortality					
	New cases			Rates			Deaths			Rates		
	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons
0–4	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
5–9	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
10–14	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
15–19	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
20–24	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
25–29	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
30–34	0	0	0	0.0	0.0	0.0	1	0	1	1.8	0.0	0.9
35–39	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
40–44	0	1	1	0.0	1.7	0.9	0	0	0	0.0	0.0	0.0
45–49	1	2	3	1.9	3.7	2.8	1	1	2	1.9	1.8	1.8
50–54	1	1	2	1.9	1.9	1.9	1	1	2	1.9	1.9	1.9
55–59	3	1	4	6.8	2.3	4.5	1	2	3	2.3	4.5	3.4
60–64	1	1	2	2.9	2.9	2.9	2	2	4	5.9	5.8	5.8
65–69	3	3	6	10.4	9.9	10.2	2	2	4	6.9	6.6	6.8
70–74	2	4	6	7.5	13.5	10.7	2	2	4	7.5	6.7	7.1
75–79	4	4	8	18.5	14.4	16.2	2	3	5	9.3	10.8	10.1
80–84	4	8	12	31.1	39.8	36.4	4	6	10	31.1	29.8	30.3
85+	3	11	14	37.4	60.2	53.2	3	11	14	37.4	60.2	53.2
Total/Crude rate	22	36	58	2.9	4.7	3.8	19	30	49	2.5	3.9	3.2
AS Rate (Aust)				2.9	3.4	3.2				2.5	2.8	2.7
Confidence Interval (95%)				1.7 – 4.1	2.3 – 4.6	2.4 – 4.1				1.4 – 3.7	1.8 – 3.8	2.0 – 3.5
AS Rate (World)				1.9	2.2	2.1				1.7	1.8	1.8
Confidence Interval (95%)				1.2 – 2.6	1.5 – 2.9	1.6 – 2.6				1.0 – 2.4	1.1 – 2.4	1.3 – 2.2
Risk				1 in 636	1 in 560	1 in 593				1 in 709	1 in 733	1 in 721
PYLL										155	130	285
% of all cancers	0.5	1.0	0.8				0.5	1.0	0.8	1.0	2.0	1.5
Trend 1998–2002 (per annum)				-1.4%	-7.6%	-5.3%				-3.1%	1.4%	-0.6%

Table A36 – Cancer of the Pancreas (157)

South Australia Year – 2002	Incidence						Mortality					
	New cases			Rates			Deaths			Rates		
	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons
0–4	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
5–9	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
10–14	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
15–19	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
20–24	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
25–29	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
30–34	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
35–39	1	0	1	1.8	0.0	0.9	2	0	2	3.6	0.0	1.8
40–44	0	0	0	0.0	0.0	0.0	1	1	2	1.7	1.7	1.7
45–49	3	4	7	5.6	7.3	6.5	0	4	4	0.0	7.3	3.7
50–54	7	4	11	13.6	7.6	10.5	6	2	8	11.6	3.8	7.7
55–59	3	2	5	6.8	4.5	5.6	5	4	9	11.3	9.0	10.2
60–64	6	4	10	17.6	11.5	14.5	7	2	9	20.5	5.8	13.1
65–69	5	9	14	17.4	29.7	23.7	16	12	28	55.5	39.6	47.4
70–74	11	8	19	41.4	26.9	33.8	12	7	19	45.2	23.5	33.8
75–79	15	16	31	69.5	57.5	62.7	10	16	26	46.3	57.5	52.6
80–84	13	9	22	101.2	44.7	66.7	8	16	24	62.3	79.5	72.8
85+	5	15	20	62.3	82.1	76.0	4	13	17	49.9	71.1	64.6
Total/Crude rate	69	71	140	9.2	9.3	9.2	71	77	148	9.5	10.0	9.7
AS Rate (Aust)				8.9	7.1	7.9				9.1	7.6	8.4
Confidence Interval (95%)				6.8 – 11.1	5.4 – 8.8	6.6 – 9.2				6.9 – 11.2	5.9 – 9.4	7.1 – 9.8
AS Rate (World)				5.9	4.7	5.3				6.5	5.1	5.8
Confidence Interval (95%)				4.7 – 7.2	3.6 – 5.8	4.4 – 6.1				5.1 – 7.9	4.0 – 6.2	4.9 – 6.7
Risk				1 in 193	1 in 229	1 in 210				1 in 134	1 in 221	1 in 168
PYLL										568	390	958
% of all cancers	1.7	2.0	1.8				1.7	2.0	1.8	3.9	5.1	4.5
Trend 1998–2002 (per annum)				-0.9%	-3.1%	-1.7%				-0.5%	2.1%	1.2%

Note: All rates are expressed per 100,000 population. Age-standardised rates (AS Rate) use the Australian 2001 and New World Standard populations (see Methods)

Table A37 – Cancer of the ill-defined sites - digestive organs (159)

South Australia Year – 2002	Incidence						Mortality					
	New cases			Rates			Deaths			Rates		
	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons
0-4	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
5-9	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
10-14	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
15-19	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
20-24	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
25-29	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
30-34	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
35-39	0	1	1	0.0	1.8	0.9	0	0	0	0.0	0.0	0.0
40-44	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
45-49	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
50-54	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
55-59	1	2	3	2.3	4.5	3.4	1	0	1	2.3	0.0	1.1
60-64	0	1	1	0.0	2.9	1.5	0	1	1	0.0	2.9	1.5
65-69	0	0	0	0.0	0.0	0.0	1	0	1	3.5	0.0	1.7
70-74	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
75-79	0	1	1	0.0	3.6	2.0	0	1	1	0.0	3.6	2.0
80-84	0	4	4	0.0	19.9	12.1	0	3	3	0.0	14.9	9.1
85+	1	2	3	12.5	10.9	11.4	1	2	3	12.5	10.9	11.4
Total/Crude rate	2	11	13	0.3	1.4	0.9	3	7	10	0.4	0.9	0.7
AS Rate (Aust)				0.3	1.1	0.7				0.4	0.6	0.5
Confidence Interval (95%)				0.0 – 0.7	0.4 – 1.7	0.3 – 1.1				0.0 – 0.9	0.2 – 1.1	0.2 – 0.9
AS Rate (World)				0.2	0.7	0.5				0.3	0.4	0.3
Confidence Interval (95%)				0.0 – 0.4	0.3 – 1.2	0.2 – 0.7				0.0 – 0.6	0.1 – 0.6	0.1 – 0.5
Risk				1 in 8,821	1 in 2,178	1 in 3,485				1 in 3,486	1 in 6,943	1 in 4,679
PYLL										25	13	38
% of all cancers	0.0	0.3	0.2				0.0	0.3	0.2	0.2	0.5	0.3
Trend 1998-2002 (per annum)				-11.1%	-3.3%	-7.5%				-9.8%	-14.7%	-13.2%

Table A38 – Cancer of the Digestive organs (150-159)

South Australia Year – 2002	Incidence						Mortality					
	New cases			Rates			Deaths			Rates		
	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons
0-4	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
5-9	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
10-14	1	1	2	1.9	2.0	2.0	0	1	1	0.0	2.0	1.0
15-19	0	2	2	0.0	4.0	1.9	0	0	0	0.0	0.0	0.0
20-24	1	0	1	2.0	0.0	1.0	0	0	0	0.0	0.0	0.0
25-29	2	1	3	4.0	2.1	3.1	1	0	1	2.0	0.0	1.0
30-34	3	1	4	5.4	1.9	3.7	3	1	4	5.4	1.9	3.7
35-39	10	8	18	17.9	14.4	16.2	3	2	5	5.4	3.6	4.5
40-44	13	14	27	22.3	23.8	23.0	4	4	8	6.9	6.8	6.8
45-49	30	22	52	55.9	40.2	48.0	11	15	26	20.5	27.4	24.0
50-54	67	37	104	129.8	70.1	99.6	31	10	41	60.1	18.9	39.3
55-59	77	65	142	174.6	146.3	160.4	44	25	69	99.8	56.3	77.9
60-64	114	52	166	334.5	149.8	241.3	46	25	71	135.0	72.0	103.2
65-69	129	81	210	447.7	267.4	355.3	71	34	105	246.4	112.3	177.7
70-74	175	112	287	659.0	376.8	509.9	76	45	121	286.2	151.4	215.0
75-79	160	145	305	741.3	520.8	617.1	83	72	155	384.5	258.6	313.6
80-84	96	111	207	747.1	551.6	627.7	63	74	137	490.3	367.7	415.5
85+	68	140	208	847.6	765.9	790.8	61	106	167	760.3	579.9	634.9
Total/Crude rate	946	792	1,738	125.9	103.2	114.4	497	414	911	66.2	53.9	60.0
AS Rate (Aust)				121.4	80.7	99.7				64.9	40.3	51.5
Confidence Interval (95%)				113.6 – 129.2	74.9 – 86.4	95.0 – 104.4				59.2 – 70.7	36.3 – 44.2	48.1 – 54.8
AS Rate (World)				85.5	55.6	69.7				43.9	26.3	34.5
Confidence Interval (95%)				80.4 – 90.5	51.7 – 59.4	66.6 – 72.9				40.4 – 47.4	23.8 – 28.8	32.4 – 36.6
Risk				1 in 11	1 in 19	1 in 14				1 in 24	1 in 45	1 in 31
PYLL										3,485	2,065	5,550
% of all cancers	23.1	21.9	22.5				23.1	21.9	22.5	27.3	27.5	27.4
Trend 1998-2002 (per annum)				-0.2%	-0.3%	-0.1%				-0.3%	-1.5%	-0.7%

Table A39 – Cancer of the Nose,nasal cavities,middle ear & accessory sinuses (160)

South Australia Year – 2002	Incidence						Mortality					
	New cases			Rates			Deaths			Rates		
	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons
0-4	0	0	0	0.0	0.0	0.0	1	0	1	2.2	0.0	1.1
5-9	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
10-14	1	0	1	1.9	0.0	1.0	0	0	0	0.0	0.0	0.0
15-19	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
20-24	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
25-29	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
30-34	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
35-39	1	0	1	1.8	0.0	0.9	0	0	0	0.0	0.0	0.0
40-44	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
45-49	1	0	1	1.9	0.0	0.9	0	0	0	0.0	0.0	0.0
50-54	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
55-59	2	1	3	4.5	2.3	3.4	0	0	0	0.0	0.0	0.0
60-64	0	1	1	0.0	2.9	1.5	0	0	0	0.0	0.0	0.0
65-69	0	0	0	0.0	0.0	0.0	1	0	1	3.5	0.0	1.7
70-74	1	1	2	3.8	3.4	3.6	1	0	1	3.8	0.0	1.8
75-79	0	0	0	0.0	0.0	0.0	1	0	1	4.6	0.0	2.0
80-84	1	0	1	7.8	0.0	3.0	1	0	1	7.8	0.0	3.0
85+	1	0	1	12.5	0.0	3.8	1	0	1	12.5	0.0	3.8
Total/Crude rate	8	3	11	1.1	0.4	0.7	6	0	6	0.8	0.0	0.4
AS Rate (Aust)				1.1	0.3	0.7				0.8	0.0	0.3
Confidence Interval (95%)				0.3 – 1.8	0.0 – 0.7	0.3 – 1.1				0.2 – 1.5	0.0 – 0.0	0.1 – 0.6
AS Rate (World)				0.8	0.3	0.5				0.6	0.0	0.3
Confidence Interval (95%)				0.3 – 1.4	0.0 – 0.6	0.2 – 0.9				0.1 – 1.1	0.0 – 0.0	0.0 – 0.5
Risk				1 in 1,441	1 in 2,354	1 in 1,785				1 in 2,129	N/A	1 in 4,376
PYLL										83	0	83
% of all cancers	0.2	0.1	0.1				0.2	0.1	0.1	0.3	0.0	0.2
Trend 1998-2002 (per annum)				1.8%	-2.9%	-1.9%				23.9%	-27.1%	-1.5%

Note: All rates are expressed per 100,000 population. Age-standardised rates (AS Rate) use the Australian 2001 and New World Standard populations (see Methods)

Table A40 – Cancer of the Larynx (161)

South Australia Year – 2002	Incidence						Mortality					
	New cases			Rates			Deaths			Rates		
	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons
0–4	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
5–9	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
10–14	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
15–19	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
20–24	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
25–29	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
30–34	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
35–39	0	1	1	0.0	1.8	0.9	1	0	1	1.8	0.0	0.9
40–44	1	0	1	1.7	0.0	0.9	0	0	0	0.0	0.0	0.0
45–49	4	0	4	7.5	0.0	3.7	0	0	0	0.0	0.0	0.0
50–54	4	1	5	7.8	1.9	4.8	1	0	1	1.9	0.0	1.0
55–59	6	1	7	13.6	2.3	7.9	0	0	0	0.0	0.0	0.0
60–64	4	1	5	11.7	2.9	7.3	0	0	0	0.0	0.0	0.0
65–69	3	1	4	10.4	3.3	6.8	4	0	4	13.9	0.0	6.8
70–74	5	1	6	18.8	3.4	10.7	6	0	6	22.6	0.0	10.7
75–79	8	2	10	37.1	7.2	20.2	3	0	3	13.9	0.0	6.1
80–84	2	0	2	15.6	0.0	6.1	5	0	5	38.9	0.0	15.2
85+	0	0	0	0.0	0.0	0.0	1	0	1	12.5	0.0	3.8
Total/Crude rate	37	8	45	4.9	1.0	3.0	21	0	21	2.8	0.0	1.4
AS Rate (Aust)				4.6	0.9	2.7				2.7	0.0	1.2
Confidence Interval (95%)				3.1 – 6.1	0.3 – 1.6	1.9 – 3.4				1.5 – 3.9	0.0 – 0.0	0.7 – 1.7
AS Rate (World)				3.5	0.7	2.0				1.8	0.0	0.8
Confidence Interval (95%)				2.4 – 4.5	0.2 – 1.2	1.5 – 2.6				1.1 – 2.4	0.0 – 0.0	0.5 – 1.1
Risk				1 in 280	1 in 1,291	1 in 467				1 in 498	N/A	1 in 1,038
PYLL										105	0	105
% of all cancers	0.9	0.2	0.6				0.9	0.2	0.6	1.2	0.0	0.6
Trend 1998–2002 (per annum)				-7.8%	27.7%	-4.1%				-1.0%	-16.3%	-0.7%

Table A41 – Cancer of the Trachea,bronchus & lung (162)

South Australia Year – 2002	Incidence						Mortality					
	New cases			Rates			Deaths			Rates		
	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons
0–4	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
5–9	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
10–14	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
15–19	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
20–24	1	0	1	2.0	0.0	1.0	0	0	0	0.0	0.0	0.0
25–29	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
30–34	0	2	2	0.0	3.7	1.8	0	1	1	0.0	1.9	0.9
35–39	0	0	0	0.0	0.0	0.0	1	0	1	1.8	0.0	0.9
40–44	8	5	13	13.7	8.5	11.1	2	4	6	3.4	6.8	5.1
45–49	9	7	16	16.8	12.8	14.8	2	7	9	3.7	12.8	8.3
50–54	17	19	36	32.9	36.0	34.5	18	10	28	34.9	18.9	26.8
55–59	27	16	43	61.2	36.0	48.6	29	10	39	65.8	22.5	44.1
60–64	33	31	64	96.8	89.3	93.0	31	16	47	91.0	46.1	68.3
65–69	66	34	100	229.0	112.3	169.2	54	31	85	187.4	102.4	143.8
70–74	106	53	159	399.1	178.3	282.5	79	32	111	297.5	107.7	197.2
75–79	87	37	124	403.1	132.9	250.9	98	34	132	454.0	122.1	267.1
80–84	56	34	90	435.8	168.9	272.9	51	30	81	396.9	149.1	245.6
85+	33	30	63	411.3	164.1	239.5	33	33	66	411.3	180.5	250.9
Total/Crude rate	443	268	711	59.0	34.9	46.8	398	208	606	53.0	27.1	39.9
AS Rate (Aust)				56.8	28.4	40.7				51.1	21.4	34.4
Confidence Interval (95%)				51.5 – 62.2	24.9 – 31.8	37.7 – 43.7				46.1 – 56.2	18.4 – 24.3	31.6 – 37.1
AS Rate (World)				38.3	20.3	28.3				33.9	14.8	23.3
Confidence Interval (95%)				35.1 – 41.6	17.9 – 22.7	26.3 – 30.3				30.9 – 36.9	12.8 – 16.8	21.6 – 25.1
Risk				1 in 24	1 in 42	1 in 31				1 in 30	1 in 63	1 in 41
PYLL										2,060	1,278	3,338
% of all cancers	10.8	7.4	9.2				10.8	7.4	9.2	21.9	13.8	18.2
Trend 1998–2002 (per annum)				-4.0%	6.9%	-0.6%				-4.1%	6.0%	-1.1%

Table A42 – Cancer of the Pleura (163)

South Australia Year – 2002	Incidence						Mortality					
	New cases			Rates			Deaths			Rates		
	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons
0–4	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
5–9	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
10–14	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
15–19	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
20–24	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
25–29	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
30–34	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
35–39	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
40–44	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
45–49	0	1	1	0.0	1.8	0.9	0	1	1	0.0	1.8	0.9
50–54	1	1	2	1.9	1.9	1.9	1	0	1	1.9	0.0	1.0
55–59	5	0	5	11.3	0.0	5.6	5	1	6	11.3	2.3	6.8
60–64	5	0	5	14.7	0.0	7.3	2	2	4	5.9	5.8	5.8
65–69	9	2	11	31.2	6.6	18.6	9	0	9	31.2	0.0	15.2
70–74	12	0	12	45.2	0.0	21.3	8	1	9	30.1	3.4	16.0
75–79	7	2	9	32.4	7.2	18.2	10	0	10	46.3	0.0	20.2
80–84	5	0	5	38.9	0.0	15.2	6	0	6	46.7	0.0	18.2
85+	3	2	5	37.4	10.9	19.0	1	0	1	12.5	0.0	3.8
Total/Crude rate	47	8	55	6.3	1.0	3.6	42	5	47	5.6	0.7	3.1
AS Rate (Aust)				6.0	0.8	3.2				5.3	0.6	2.7
Confidence Interval (95%)				4.3 – 7.7	0.2 – 1.4	2.3 – 4.0				3.7 – 6.9	0.1 – 1.1	1.9 – 3.5
AS Rate (World)				4.2	0.6	2.2				3.6	0.5	1.9
Confidence Interval (95%)				3.1 – 5.3	0.2 – 1.0	1.7 – 2.8				2.6 – 4.6	0.1 – 0.9	1.4 – 2.5
Risk				1 in 192	1 in 1,937	1 in 360				1 in 249	1 in 1,515	1 in 438
PYLL										223	73	295
% of all cancers	1.1	0.2	0.7				1.1	0.2	0.7	2.3	0.3	1.4
Trend 1998–2002 (per annum)				1.4%	15.0%	4.3%				11.3%	11.4%	12.4%

Note: All rates are expressed per 100,000 population. Age-standardised rates (AS Rate) use the Australian 2001 and New World Standard populations (see Methods)

Table A43 – Cancer of the Respiratory organs (160-163)

South Australia Year – 2002	Incidence						Mortality					
	New cases			Rates			Deaths			Rates		
	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons
0-4	0	0	0	0.0	0.0	0.0	1	0	1	2.2	0.0	1.1
5-9	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
10-14	1	0	1	1.9	0.0	1.0	0	0	0	0.0	0.0	0.0
15-19	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
20-24	1	0	1	2.0	0.0	1.0	0	0	0	0.0	0.0	0.0
25-29	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
30-34	0	2	2	0.0	3.7	1.8	0	1	1	0.0	1.9	0.9
35-39	1	1	2	1.8	1.8	1.8	2	0	2	3.6	0.0	1.8
40-44	9	5	14	15.4	8.5	11.9	2	4	6	3.4	6.8	5.1
45-49	14	8	22	26.1	14.6	20.3	2	8	10	3.7	14.6	9.2
50-54	22	21	43	42.6	39.8	41.2	20	10	30	38.8	18.9	28.7
55-59	40	18	58	90.7	40.5	65.5	34	11	45	77.1	24.8	50.8
60-64	42	33	75	123.2	95.1	109.0	33	18	51	96.8	51.9	74.1
65-69	78	37	115	270.7	122.2	194.6	68	31	99	236.0	102.4	167.5
70-74	124	55	179	466.9	185.0	318.0	94	33	127	354.0	111.0	225.7
75-79	102	41	143	472.6	147.3	289.3	112	34	146	518.9	122.1	295.4
80-84	64	34	98	498.1	168.9	297.2	63	30	93	490.3	149.1	282.0
85+	37	32	69	461.2	175.1	262.3	36	33	69	448.7	180.5	262.3
Total/Crude rate	535	287	822	71.2	37.4	54.1	467	213	680	62.2	27.8	44.8
AS Rate (Aust)				68.5	30.5	47.2				59.9	21.9	38.6
Confidence Interval (95%)				62.6 – 74.3	26.9 – 34.0	44.0 – 50.4				54.5 – 65.4	19.0 – 24.9	35.7 – 41.5
AS Rate (World)				46.8	21.9	33.1				39.9	15.3	26.4
Confidence Interval (95%)				43.2 – 50.5	19.4 – 24.4	31.0 – 35.3				36.6 – 43.1	13.3 – 17.4	24.5 – 28.2
Risk				1 in 20	1 in 40	1 in 27				1 in 25	1 in 61	1 in 36
PYLL										2,470	1,350	3,820
% of all cancers	13.1	7.9	10.7				13.1	7.9	10.7	25.6	14.2	20.5
Trend 1998-2002 (per annum)				-3.8%	7.4%	-0.5%				-3.0%	5.6%	-0.4%

Table A44 – Cancer of the Thymus, heart and mediastinum (164)

South Australia Year – 2002	Incidence						Mortality					
	New cases			Rates			Deaths			Rates		
	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons
0-4	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
5-9	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
10-14	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
15-19	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
20-24	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
25-29	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
30-34	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
35-39	0	0	0	0.0	0.0	0.0	1	0	1	1.8	0.0	0.9
40-44	1	0	1	1.7	0.0	0.9	0	0	0	0.0	0.0	0.0
45-49	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
50-54	1	0	1	1.9	0.0	1.0	0	0	0	0.0	0.0	0.0
55-59	3	0	3	6.8	0.0	3.4	1	0	1	2.3	0.0	1.1
60-64	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
65-69	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
70-74	0	1	1	0.0	3.4	1.8	0	0	0	0.0	0.0	0.0
75-79	0	1	1	0.0	3.6	2.0	0	0	0	0.0	0.0	0.0
80-84	0	0	0	0.0	0.0	0.0	1	0	1	7.8	0.0	3.0
85+	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
Total/Crude rate	5	2	7	0.7	0.3	0.5	3	0	3	0.4	0.0	0.2
AS Rate (Aust)				0.6	0.2	0.4				0.4	0.0	0.2
Confidence Interval (95%)				0.1 – 1.2	0.0 – 0.5	0.1 – 0.7				0.0 – 0.8	0.0 – 0.0	0.0 – 0.4
AS Rate (World)				0.5	0.1	0.3				0.3	0.0	0.1
Confidence Interval (95%)				0.1 – 0.9	0.0 – 0.3	0.1 – 0.6				0.0 – 0.6	0.0 – 0.0	0.0 – 0.3
Risk				1 in 1,913	1 in 5,945	1 in 2,867				1 in 4,933	N/A	1 in 9,865
PYLL										55	0	55
% of all cancers	0.1	0.1	0.1				0.1	0.1	0.1	0.2	0.0	0.1
Trend 1998-2002 (per annum)				-0.2%	0.0%	0.4%				38.0%	-31.9%	-11.2%

Table A45 – Cancer of the Bone and articular cartilage (170)

South Australia Year – 2002	Incidence						Mortality					
	New cases			Rates			Deaths			Rates		
	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons
0-4	1	0	1	2.2	0.0	1.1	0	0	0	0.0	0.0	0.0
5-9	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
10-14	0	1	1	0.0	2.0	1.0	1	0	1	1.9	0.0	1.0
15-19	3	0	3	5.7	0.0	2.9	0	0	0	0.0	0.0	0.0
20-24	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
25-29	1	0	1	2.0	0.0	1.0	0	0	0	0.0	0.0	0.0
30-34	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
35-39	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
40-44	1	0	1	1.7	0.0	0.9	0	0	0	0.0	0.0	0.0
45-49	1	0	1	1.9	0.0	0.9	0	0	0	0.0	0.0	0.0
50-54	2	0	2	3.9	0.0	1.9	0	0	0	0.0	0.0	0.0
55-59	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
60-64	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
65-69	1	0	1	3.5	0.0	1.7	0	0	0	0.0	0.0	0.0
70-74	1	1	2	3.8	3.4	3.6	1	0	1	3.8	0.0	1.8
75-79	1	1	2	4.6	3.6	4.0	0	1	1	0.0	3.6	2.0
80-84	1	0	1	7.8	0.0	3.0	1	0	1	7.8	0.0	3.0
85+	1	0	1	12.5	0.0	3.8	0	0	0	0.0	0.0	0.0
Total/Crude rate	14	3	17	1.9	0.4	1.1	3	1	4	0.4	0.1	0.3
AS Rate (Aust)				1.9	0.3	1.1				0.4	0.1	0.2
Confidence Interval (95%)				0.9 – 2.9	0.0 – 0.7	0.6 – 1.6				0.0 – 0.8	0.0 – 0.3	0.0 – 0.5
AS Rate (World)				1.7	0.3	1.0				0.3	0.1	0.2
Confidence Interval (95%)				0.8 – 2.6	0.0 – 0.7	0.5 – 1.5				0.0 – 0.7	0.0 – 0.1	0.0 – 0.4
Risk				1 in 816	1 in 3,698	1 in 1,337				1 in 3,512	N/A	1 in 7,221
PYLL										65	0	65
% of all cancers	0.3	0.1	0.2				0.3	0.1	0.2	0.2	0.1	0.1
Trend 1998-2002 (per annum)				32.3%	-20.8%	-3.4%				7.7%	-19.5%	-7.1%

Note: All rates are expressed per 100,000 population. Age-standardised rates (AS Rate) use the Australian 2001 and New World Standard populations (see Methods)

Table A46 – Cancer of the Connective & other soft tissue (171)

South Australia Year – 2002	Incidence						Mortality					
	New cases			Rates			Deaths			Rates		
	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons
0-4	1	0	1	2.2	0.0	1.1	1	0	1	2.2	0.0	1.1
5-9	0	2	2	0.0	4.2	2.0	0	1	1	0.0	2.1	1.0
10-14	0	1	1	0.0	2.0	1.0	0	0	0	0.0	0.0	0.0
15-19	3	0	3	5.7	0.0	2.9	1	0	1	1.9	0.0	1.0
20-24	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
25-29	1	0	1	2.0	0.0	1.0	1	1	2	2.0	2.1	2.1
30-34	2	1	3	3.6	1.9	2.7	0	0	0	0.0	0.0	0.0
35-39	5	2	7	8.9	3.6	6.3	0	0	0	0.0	0.0	0.0
40-44	0	1	1	0.0	1.7	0.9	0	1	1	0.0	1.7	0.9
45-49	1	3	4	1.9	5.5	3.7	0	1	1	0.0	1.8	0.9
50-54	4	2	6	7.8	3.8	5.7	1	0	1	1.9	0.0	1.0
55-59	3	2	5	6.8	4.5	5.6	0	3	3	0.0	6.8	3.4
60-64	3	0	3	8.8	0.0	4.4	2	0	2	5.9	0.0	2.9
65-69	5	0	5	17.4	0.0	8.5	4	2	6	13.9	6.6	10.2
70-74	3	1	4	11.3	3.4	7.1	2	1	3	7.5	3.4	5.3
75-79	7	3	10	32.4	10.8	20.2	1	1	2	4.6	3.6	4.0
80-84	5	3	8	38.9	14.9	24.3	6	2	8	46.7	9.9	24.3
85+	2	1	3	24.9	5.5	11.4	1	0	1	12.5	0.0	3.8
Total/Crude rate	45	22	67	6.0	2.9	4.4	20	13	33	2.7	1.7	2.2
AS Rate (Aust)				5.9	2.6	4.1				2.6	1.5	2.0
Confidence Interval (95%)				4.2 – 7.6	1.5 – 3.7	3.1 – 5.1				1.5 – 3.8	0.7 – 2.3	1.3 – 2.6
AS Rate (World)				4.7	2.2	3.4				2.0	1.3	1.6
Confidence Interval (95%)				3.4 – 6.0	1.2 – 3.2	2.5 – 4.2				1.1 – 2.9	0.5 – 2.0	1.0 – 2.1
Risk				1 in 263	1 in 656	1 in 378				1 in 567	1 in 819	1 in 675
PYLL										260	245	505
% of all cancers	1.1	0.6	0.9				1.1	0.6	0.9	1.1	0.9	1.0
Trend 1998–2002 (per annum)				-3.6%	-8.6%	-5.8%				-2.3%	17.3%	3.9%

Table A47 – Melanoma of skin (172)

South Australia Year – 2002	Incidence						Mortality					
	New cases			Rates			Deaths			Rates		
	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons
0-4	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
5-9	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
10-14	0	1	1	0.0	2.0	1.0	0	0	0	0.0	0.0	0.0
15-19	0	3	3	0.0	5.9	2.9	0	0	0	0.0	0.0	0.0
20-24	2	11	13	4.0	23.4	13.4	0	0	0	0.0	0.0	0.0
25-29	7	14	21	14.1	29.5	21.7	0	1	1	0.0	2.1	1.0
30-34	10	20	30	18.1	37.1	27.5	1	0	1	1.8	0.0	0.9
35-39	15	15	30	26.8	27.1	26.9	3	0	3	5.4	0.0	2.7
40-44	30	21	51	51.5	35.7	43.5	2	3	5	3.4	5.1	4.3
45-49	30	24	54	55.9	43.9	49.8	3	0	3	5.6	0.0	2.8
50-54	31	38	69	60.1	72.0	66.1	0	1	1	0.0	1.9	1.0
55-59	28	23	51	63.5	51.8	57.6	3	1	4	6.8	2.3	4.5
60-64	36	27	63	105.6	77.8	91.6	10	0	10	29.3	0.0	14.5
65-69	26	24	50	90.2	79.2	84.6	7	3	10	24.3	9.9	16.9
70-74	42	30	72	158.2	100.9	127.9	7	1	8	26.4	3.4	14.2
75-79	44	28	72	203.9	100.6	145.7	7	2	9	32.4	7.2	18.2
80-84	21	32	53	163.4	159.0	160.7	5	6	11	38.9	29.8	33.4
85+	18	19	37	224.4	103.9	140.7	3	5	8	37.4	27.4	30.4
Total/Crude rate	340	330	670	45.3	43.0	44.1	51	23	74	6.8	3.0	4.9
AS Rate (Aust)				44.0	38.7	40.7				6.6	2.3	4.3
Confidence Interval (95%)				39.3 – 48.7	34.4 – 42.9	37.6 – 43.8				4.8 – 8.4	1.3 – 3.3	3.3 – 5.3
AS Rate (World)				33.6	32.0	32.5				4.9	1.6	3.2
Confidence Interval (95%)				30.3 – 37.0	28.5 – 35.4	30.1 – 34.9				3.6 – 6.1	0.9 – 2.3	2.5 – 3.9
Risk				1 in 31	1 in 35	1 in 33				1 in 195	1 in 813	1 in 319
PYLL										550	210	760
% of all cancers	8.3	9.1	8.7				8.3	9.1	8.7	2.8	1.5	2.2
Trend 1998–2002 (per annum)				2.0%	1.1%	1.8%				3.3%	-5.2%	0.5%

Table A48 – Melanoma of lip (172.0)

South Australia Year – 2002	Incidence						Mortality					
	New cases			Rates			Deaths			Rates		
	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons
0-4	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
5-9	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
10-14	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
15-19	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
20-24	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
25-29	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
30-34	0	1	1	0.0	1.9	0.9	0	0	0	0.0	0.0	0.0
35-39	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
40-44	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
45-49	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
50-54	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
55-59	1	0	1	2.3	0.0	1.1	0	0	0	0.0	0.0	0.0
60-64	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
65-69	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
70-74	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
75-79	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
80-84	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
85+	1	0	1	12.5	0.0	3.8	0	0	0	0.0	0.0	0.0
Total/Crude rate	2	1	3	0.3	0.1	0.2	0	0	0	0.0	0.0	0.0
AS Rate (Aust)				0.3	0.1	0.2				0.0	0.0	0.0
Confidence Interval (95%)				0.0 – 0.7	0.0 – 0.4	0.0 – 0.4				0.0 – 0.0	0.0 – 0.0	0.0 – 0.0
AS Rate (World)				0.2	0.1	0.1				0.0	0.0	0.0
Confidence Interval (95%)				0.0 – 0.4	0.0 – 0.4	0.0 – 0.3				0.0 – 0.0	0.0 – 0.0	0.0 – 0.0
Risk				1 in 8,821	1 in 10,796	1 in 9,775				N/A	N/A	N/A
PYLL										0	0	0
% of all cancers	0.0	0.0	0.0				0.0	0.0	0.0	0.0	0.0	0.0
Trend 1998–2002 (per annum)				undefined	27.7%	202.8%				0.0%	0.0%	0.0%

Note: All rates are expressed per 100,000 population. Age-standardised rates (AS Rate) use the Australian 2001 and New World Standard populations (see Methods)

Table A49 – Melanoma of eyelid, including canthus (172.1)

South Australia Year – 2002	Incidence						Mortality					
	New cases			Rates			Deaths			Rates		
	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons
0–4	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
5–9	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
10–14	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
15–19	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
20–24	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
25–29	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
30–34	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
35–39	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
40–44	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
45–49	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
50–54	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
55–59	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
60–64	1	0	1	2.9	0.0	1.5	0	0	0	0.0	0.0	0.0
65–69	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
70–74	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
75–79	1	0	1	4.6	0.0	2.0	0	0	0	0.0	0.0	0.0
80–84	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
85+	0	1	1	0.0	5.5	3.8	0	0	0	0.0	0.0	0.0
Total/Crude rate	2	1	3	0.3	0.1	0.2	0	0	0	0.0	0.0	0.0
AS Rate (Aust)				0.2	0.1	0.2				0.0	0.0	0.0
Confidence Interval (95%)				0.0 – 0.6	0.0 – 0.2	0.0 – 0.4				0.0 – 0.0	0.0 – 0.0	0.0 – 0.0
AS Rate (World)				0.2	0.0	0.1				0.0	0.0	0.0
Confidence Interval (95%)				0.0 – 0.4	0.0 – 0.1	0.0 – 0.2				0.0 – 0.0	0.0 – 0.0	0.0 – 0.0
Risk				1 in 6,817	N/A	1 in 13,759				N/A	N/A	N/A
PYLL										0	0	0
% of all cancers	0.0	0.0	0.0				0.0	0.0	0.0	0.0	0.0	0.0
Trend 1998–2002 (per annum)				6402.0%	-10.2%	23.4%				-1.3%	0.0%	-1.1%

Table A50 – Melanoma of ear (172.2)

South Australia Year – 2002	Incidence						Mortality					
	New cases			Rates			Deaths			Rates		
	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons
0–4	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
5–9	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
10–14	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
15–19	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
20–24	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
25–29	1	1	2	2.0	2.1	2.1	0	0	0	0.0	0.0	0.0
30–34	0	1	1	0.0	1.9	0.9	0	0	0	0.0	0.0	0.0
35–39	1	0	1	1.8	0.0	0.9	0	0	0	0.0	0.0	0.0
40–44	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
45–49	2	1	3	3.7	1.8	2.8	0	0	0	0.0	0.0	0.0
50–54	0	1	1	0.0	1.9	1.0	0	0	0	0.0	0.0	0.0
55–59	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
60–64	0	1	1	0.0	2.9	1.5	0	0	0	0.0	0.0	0.0
65–69	2	0	2	6.9	0.0	3.4	0	0	0	0.0	0.0	0.0
70–74	2	0	2	7.5	0.0	3.6	0	0	0	0.0	0.0	0.0
75–79	3	0	3	13.9	0.0	6.1	0	0	0	0.0	0.0	0.0
80–84	2	1	3	15.6	5.0	9.1	0	0	0	0.0	0.0	0.0
85+	1	0	1	12.5	0.0	3.8	0	0	0	0.0	0.0	0.0
Total/Crude rate	14	6	20	1.9	0.8	1.3	0	0	0	0.0	0.0	0.0
AS Rate (Aust)				1.8	0.8	1.2				0.0	0.0	0.0
Confidence Interval (95%)				0.9 – 2.8	0.1 – 1.4	0.7 – 1.7				0.0 – 0.0	0.0 – 0.0	0.0 – 0.0
AS Rate (World)				1.3	0.7	0.9				0.0	0.0	0.0
Confidence Interval (95%)				0.7 – 2.0	0.1 – 1.2	0.5 – 1.4				0.0 – 0.0	0.0 – 0.0	0.0 – 0.0
Risk				1 in 909	1 in 1,894	1 in 1,251				N/A	N/A	N/A
PYLL										0	0	0
% of all cancers	0.3	0.2	0.3				0.3	0.2	0.3	0.0	0.0	0.0
Trend 1998–2002 (per annum)				5.6%	13.5%	6.9%				-24.5%	-18.7%	-22.8%

Table A51 – Melanoma of unspecified parts of face (172.3)

South Australia Year – 2002	Incidence						Mortality					
	New cases			Rates			Deaths			Rates		
	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons
0–4	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
5–9	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
10–14	0	1	1	0.0	2.0	1.0	0	0	0	0.0	0.0	0.0
15–19	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
20–24	0	1	1	0.0	2.1	1.0	0	0	0	0.0	0.0	0.0
25–29	1	0	1	2.0	0.0	1.0	0	0	0	0.0	0.0	0.0
30–34	2	0	2	3.6	0.0	1.8	0	0	0	0.0	0.0	0.0
35–39	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
40–44	1	2	3	1.7	3.4	2.6	0	0	0	0.0	0.0	0.0
45–49	0	2	2	0.0	3.7	1.8	0	0	0	0.0	0.0	0.0
50–54	3	2	5	5.8	3.8	4.8	0	0	0	0.0	0.0	0.0
55–59	1	3	4	2.3	6.8	4.5	1	0	1	2.3	0.0	1.1
60–64	2	1	3	5.9	2.9	4.4	2	0	2	5.9	0.0	2.9
65–69	2	5	7	6.9	16.5	11.8	1	0	1	3.5	0.0	1.7
70–74	10	1	11	37.7	3.4	19.5	1	0	1	3.8	0.0	1.8
75–79	5	4	9	23.2	14.4	18.2	1	0	1	4.6	0.0	2.0
80–84	3	8	11	23.3	39.8	33.4	1	0	1	7.8	0.0	3.0
85+	1	8	9	12.5	43.8	34.2	1	1	2	12.5	5.5	7.6
Total/Crude rate	31	38	69	4.1	5.0	4.5	8	1	9	1.1	0.1	0.6
AS Rate (Aust)				4.0	3.9	4.0				1.0	0.1	0.5
Confidence Interval (95%)				2.6 – 5.4	2.6 – 5.1	3.0 – 4.9				0.3 – 1.8	0.0 – 0.2	0.2 – 0.8
AS Rate (World)				2.9	2.8	2.9				0.7	0.0	0.4
Confidence Interval (95%)				1.9 – 3.8	1.9 – 3.8	2.2 – 3.5				0.2 – 1.2	0.0 – 0.1	0.1 – 0.6
Risk				1 in 304	1 in 450	1 in 368				1 in 1,302	N/A	1 in 2,665
PYLL										53	0	53
% of all cancers	0.8	1.1	0.9				0.8	1.1	0.9	0.4	0.1	0.3
Trend 1998–2002 (per annum)				-3.1%	4.9%	0.6%				16.1%	9.1%	13.8%

Note: All rates are expressed per 100,000 population. Age-standardised rates (AS Rate) use the Australian 2001 and New World Standard populations (see Methods)

Table A52 – Melanoma of scalp and neck (172.4)

South Australia Year – 2002	Incidence						Mortality					
	New cases			Rates			Deaths			Rates		
	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons
0–4	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
5–9	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
10–14	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
15–19	0	1	1	0.0	2.0	1.0	0	0	0	0.0	0.0	0.0
20–24	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
25–29	1	0	1	2.0	0.0	1.0	0	1	1	0.0	2.1	1.0
30–34	1	0	1	1.8	0.0	0.9	0	0	0	0.0	0.0	0.0
35–39	1	0	1	1.8	0.0	0.9	0	0	0	0.0	0.0	0.0
40–44	1	1	2	1.7	1.7	1.7	0	2	2	0.0	3.4	1.7
45–49	4	1	5	7.5	1.8	4.6	2	0	2	3.7	0.0	1.8
50–54	2	0	2	3.9	0.0	1.9	0	0	0	0.0	0.0	0.0
55–59	1	1	2	2.3	2.3	2.3	1	0	1	2.3	0.0	1.1
60–64	6	2	8	17.6	5.8	11.6	1	0	1	2.9	0.0	1.5
65–69	3	1	4	10.4	3.3	6.8	0	1	1	0.0	3.3	1.7
70–74	3	0	3	11.3	0.0	5.3	3	0	3	11.3	0.0	5.3
75–79	3	0	3	13.9	0.0	6.1	2	0	2	9.3	0.0	4.0
80–84	4	3	7	31.1	14.9	21.2	0	0	0	0.0	0.0	0.0
85+	3	2	5	37.4	10.9	19.0	0	2	2	0.0	10.9	7.6
Total/Crude rate	33	12	45	4.4	1.6	3.0	9	6	15	1.2	0.8	1.0
AS Rate (Aust)				4.3	1.3	2.7				1.1	0.7	0.9
Confidence Interval (95%)				2.9 – 5.8	0.5 – 2.0	1.9 – 3.5				0.4 – 1.9	0.1 – 1.2	0.4 – 1.4
AS Rate (World)				3.2	1.0	2.0				0.8	0.6	0.7
Confidence Interval (95%)				2.2 – 4.3	0.4 – 1.6	1.4 – 2.6				0.3 – 1.3	0.1 – 1.0	0.3 – 1.0
Risk				1 in 332	1 in 1,189	1 in 526				1 in 989	1 in 2,272	1 in 1,410
PYLL										93	120	213
% of all cancers	0.8	0.3	0.6				0.8	0.3	0.6	0.5	0.4	0.5
Trend 1998–2002 (per annum)				3.1%	-5.6%	1.9%				42.1%	38.1%	46.5%

Table A53 – Melanoma of trunk (172.5)

South Australia Year – 2002	Incidence						Mortality					
	New cases			Rates			Deaths			Rates		
	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons
0–4	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
5–9	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
10–14	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
15–19	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
20–24	1	2	3	2.0	4.3	3.1	0	0	0	0.0	0.0	0.0
25–29	1	5	6	2.0	10.5	6.2	0	0	0	0.0	0.0	0.0
30–34	4	8	12	7.3	14.8	11.0	0	0	0	0.0	0.0	0.0
35–39	6	4	10	10.7	7.2	9.0	1	0	1	1.8	0.0	0.9
40–44	12	2	14	20.6	3.4	11.9	1	0	1	1.7	0.0	0.9
45–49	10	4	14	18.6	7.3	12.9	0	0	0	0.0	0.0	0.0
50–54	12	10	22	23.3	18.9	21.1	0	0	0	0.0	0.0	0.0
55–59	12	5	17	27.2	11.3	19.2	0	1	1	0.0	2.3	1.1
60–64	9	5	14	26.4	14.4	20.4	2	0	2	5.9	0.0	2.9
65–69	8	5	13	27.8	16.5	22.0	4	1	5	13.9	3.3	8.5
70–74	12	5	17	45.2	16.8	30.2	2	0	2	7.5	0.0	3.6
75–79	14	7	21	64.9	25.1	42.5	2	0	2	9.3	0.0	4.0
80–84	5	3	8	38.9	14.9	24.3	2	3	5	15.6	14.9	15.2
85+	4	1	5	49.9	5.5	19.0	0	0	0	0.0	0.0	0.0
Total/Crude rate	110	66	176	14.6	8.6	11.6	14	5	19	1.9	0.7	1.3
AS Rate (Aust)				14.2	8.1	10.8				1.8	0.5	1.1
Confidence Interval (95%)				11.5 – 16.8	6.1 – 10.1	9.2 – 12.4				0.8 – 2.7	0.1 – 0.9	0.6 – 1.6
AS Rate (World)				11.1	7.0	8.9				1.3	0.3	0.8
Confidence Interval (95%)				9.1 – 13.0	5.3 – 8.6	7.6 – 10.1				0.7 – 2.0	0.1 – 0.6	0.5 – 1.2
Risk				1 in 95	1 in 160	1 in 120				1 in 650	1 in 3,602	1 in 1,124
PYLL										130	25	155
% of all cancers	2.7	1.8	2.3				2.7	1.8	2.3	0.8	0.3	0.6
Trend 1998–2002 (per annum)				0.9%	3.6%	1.5%				-3.6%	-17.8%	-7.6%

Table A54 – Melanoma of upper limb (172.6)

South Australia Year – 2002	Incidence						Mortality					
	New cases			Rates			Deaths			Rates		
	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons
0–4	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
5–9	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
10–14	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
15–19	0	2	2	0.0	4.0	1.9	0	0	0	0.0	0.0	0.0
20–24	1	4	5	2.0	8.5	5.2	0	0	0	0.0	0.0	0.0
25–29	2	8	10	4.0	16.9	10.3	0	0	0	0.0	0.0	0.0
30–34	2	5	7	3.6	9.3	6.4	0	0	0	0.0	0.0	0.0
35–39	3	3	6	5.4	5.4	5.4	0	0	0	0.0	0.0	0.0
40–44	8	11	19	13.7	18.7	16.2	0	0	0	0.0	0.0	0.0
45–49	9	8	17	16.8	14.6	15.7	0	0	0	0.0	0.0	0.0
50–54	6	10	16	11.6	18.9	15.3	0	0	0	0.0	0.0	0.0
55–59	9	5	14	20.4	11.3	15.8	0	0	0	0.0	0.0	0.0
60–64	11	5	16	32.3	14.4	23.3	2	0	2	5.9	0.0	2.9
65–69	4	4	8	13.9	13.2	13.5	2	0	2	6.9	0.0	3.4
70–74	9	11	20	33.9	37.0	35.5	0	1	1	0.0	3.4	1.8
75–79	11	7	18	51.0	25.1	36.4	0	1	1	0.0	3.6	2.0
80–84	4	5	9	31.1	24.8	27.3	0	1	1	0.0	5.0	3.0
85+	3	4	7	37.4	21.9	26.6	1	0	1	12.5	0.0	3.8
Total/Crude rate	82	92	174	10.9	12.0	11.5	5	3	8	0.7	0.4	0.5
AS Rate (Aust)				10.5	11.2	10.7				0.7	0.3	0.5
Confidence Interval (95%)				8.3 – 12.8	8.9 – 13.5	9.1 – 12.3				0.1 – 1.2	0.0 – 0.6	0.1 – 0.8
AS Rate (World)				8.2	9.6	8.8				0.5	0.2	0.3
Confidence Interval (95%)				6.6 – 9.9	7.6 – 11.6	7.6 – 10.1				0.0 – 1.0	0.0 – 0.3	0.1 – 0.6
Risk				1 in 127	1 in 117	1 in 122				1 in 1,562	1 in 5,945	1 in 2,479
PYLL										40	3	43
% of all cancers	2.0	2.5	2.3				2.0	2.5	2.3	0.3	0.2	0.2
Trend 1998–2002 (per annum)				12.0%	6.4%	8.9%				-0.5%	-9.8%	-5.2%

Note: All rates are expressed per 100,000 population. Age-standardised rates (AS Rate) use the Australian 2001 and New World Standard populations (see Methods)

Table A55 – Melanoma of lower limb (172.7)

South Australia Year – 2002	Incidence						Mortality					
	New cases			Rates			Deaths			Rates		
	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons
0–4	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
5–9	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
10–14	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
15–19	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
20–24	0	3	3	0.0	6.4	3.1	0	0	0	0.0	0.0	0.0
25–29	1	0	1	2.0	0.0	1.0	0	0	0	0.0	0.0	0.0
30–34	0	5	5	0.0	9.3	4.6	0	0	0	0.0	0.0	0.0
35–39	4	8	12	7.2	14.4	10.8	2	0	2	3.6	0.0	1.8
40–44	6	4	10	10.3	6.8	8.5	0	1	1	0.0	1.7	0.9
45–49	4	8	12	7.5	14.6	11.1	1	0	1	1.9	0.0	0.9
50–54	8	15	23	15.5	28.4	22.0	0	1	1	0.0	1.9	1.0
55–59	4	9	13	9.1	20.3	14.7	0	0	0	0.0	0.0	0.0
60–64	4	13	17	11.7	37.4	24.7	1	0	1	2.9	0.0	1.5
65–69	6	9	15	20.8	29.7	25.4	0	1	1	0.0	3.3	1.7
70–74	6	11	17	22.6	37.0	30.2	1	0	1	3.8	0.0	1.8
75–79	7	8	15	32.4	28.7	30.3	2	1	3	9.3	3.6	6.1
80–84	3	11	14	23.3	54.7	42.5	1	1	2	7.8	5.0	6.1
85+	2	3	5	24.9	16.4	19.0	0	2	2	0.0	10.9	7.6
Total/Crude rate	55	107	162	7.3	13.9	10.7	8	7	15	1.1	0.9	1.0
AS Rate (Aust)				7.1	12.5	9.8				1.0	0.7	0.9
Confidence Interval (95%)				5.2 – 9.0	10.1 – 14.9	8.3 – 11.4				0.3 – 1.8	0.2 – 1.2	0.4 – 1.3
AS Rate (World)				5.5	10.2	7.8				0.8	0.5	0.6
Confidence Interval (95%)				4.1 – 6.8	8.3 – 12.0	6.7 – 9.0				0.3 – 1.3	0.1 – 0.8	0.3 – 0.9
Risk				1 in 188	1 in 98	1 in 129				1 in 1,648	1 in 2,901	1 in 2,116
PYLL										118	63	180
% of all cancers	1.3	3.0	2.1				1.3	3.0	2.1	0.4	0.5	0.5
Trend 1998–2002 (per annum)				-0.8%	-3.0%	-2.2%				11.8%	6.5%	9.6%

Table A56 – Melanoma of other site (172.8)

South Australia Year – 2002	Incidence						Mortality					
	New cases			Rates			Deaths			Rates		
	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons
0–4	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
5–9	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
10–14	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
15–19	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
20–24	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
25–29	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
30–34	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
35–39	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
40–44	1	0	1	1.7	0.0	0.9	0	0	0	0.0	0.0	0.0
45–49	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
50–54	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
55–59	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
60–64	1	0	1	2.9	0.0	1.5	0	0	0	0.0	0.0	0.0
65–69	1	0	1	3.5	0.0	1.7	0	0	0	0.0	0.0	0.0
70–74	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
75–79	0	1	1	0.0	3.6	2.0	0	0	0	0.0	0.0	0.0
80–84	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
85+	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
Total/Crude rate	3	1	4	0.4	0.1	0.3	0	0	0	0.0	0.0	0.0
AS Rate (Aust)				0.4	0.1	0.2				0.0	0.0	0.0
Confidence Interval (95%)				0.0 – 0.8	0.0 – 0.3	0.0 – 0.5				0.0 – 0.0	0.0 – 0.0	0.0 – 0.0
AS Rate (World)				0.3	0.1	0.2				0.0	0.0	0.0
Confidence Interval (95%)				0.0 – 0.7	0.0 – 0.1	0.0 – 0.4				0.0 – 0.0	0.0 – 0.0	0.0 – 0.0
Risk				1 in 2,463	N/A	1 in 5,002				N/A	N/A	N/A
PYLL										0	0	0
% of all cancers	0.1	0.0	0.1				0.1	0.0	0.1	0.0	0.0	0.0
Trend 1998–2002 (per annum)				0.7%	82.9%	22.7%				0.0%	0.0%	0.0%

Table A57 – Melanoma - site unspecified (172.9)

South Australia Year – 2002	Incidence						Mortality					
	New cases			Rates			Deaths			Rates		
	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons
0–4	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
5–9	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
10–14	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
15–19	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
20–24	0	1	1	0.0	2.1	1.0	0	0	0	0.0	0.0	0.0
25–29	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
30–34	1	0	1	1.8	0.0	0.9	1	0	1	1.8	0.0	0.9
35–39	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
40–44	1	1	2	1.7	1.7	1.7	1	0	1	1.7	0.0	0.9
45–49	1	0	1	1.9	0.0	0.9	0	0	0	0.0	0.0	0.0
50–54	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
55–59	0	0	0	0.0	0.0	0.0	1	0	1	2.3	0.0	1.1
60–64	2	0	2	5.9	0.0	2.9	2	0	2	5.9	0.0	2.9
65–69	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
70–74	0	2	2	0.0	6.7	3.6	0	0	0	0.0	0.0	0.0
75–79	0	1	1	0.0	3.6	2.0	0	0	0	0.0	0.0	0.0
80–84	0	1	1	0.0	5.0	3.0	1	1	2	7.8	5.0	6.1
85+	3	0	3	37.4	0.0	11.4	1	0	1	12.5	0.0	3.8
Total/Crude rate	8	6	14	1.1	0.8	0.9	7	1	8	0.9	0.1	0.5
AS Rate (Aust)				1.2	0.7	0.8				0.9	0.1	0.5
Confidence Interval (95%)				0.3 – 2.0	0.1 – 1.2	0.4 – 1.3				0.2 – 1.6	0.0 – 0.3	0.1 – 0.8
AS Rate (World)				0.8	0.5	0.6				0.7	0.0	0.4
Confidence Interval (95%)				0.3 – 1.3	0.1 – 1.0	0.3 – 1.0				0.2 – 1.2	0.0 – 0.1	0.1 – 0.6
Risk				1 in 1,776	1 in 1,896	1 in 1,812				1 in 1,715	N/A	1 in 3,445
PYLL										118	0	118
% of all cancers	0.2	0.2	0.2				0.2	0.2	0.2	0.4	0.1	0.2
Trend 1998–2002 (per annum)				-13.0%	-7.6%	-10.6%				-3.3%	-14.7%	-6.5%

Note: All rates are expressed per 100,000 population. Age-standardised rates (AS Rate) use the Australian 2001 and New World Standard populations (see Methods)

Table A58 – Cancer of other skin - lip & anus (173)

South Australia Year – 2002	Incidence						Mortality					
	New cases			Rates			Deaths			Rates		
	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons
0–4	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
5–9	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
10–14	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
15–19	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
20–24	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
25–29	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
30–34	2	0	2	3.6	0.0	1.8	0	0	0	0.0	0.0	0.0
35–39	1	2	3	1.8	3.6	2.7	0	0	0	0.0	0.0	0.0
40–44	4	0	4	6.9	0.0	3.4	0	0	0	0.0	0.0	0.0
45–49	2	1	3	3.7	1.8	2.8	0	0	0	0.0	0.0	0.0
50–54	4	2	6	7.8	3.8	5.7	0	0	0	0.0	0.0	0.0
55–59	3	4	7	6.8	9.0	7.9	0	0	0	0.0	0.0	0.0
60–64	6	2	8	17.6	5.8	11.6	0	0	0	0.0	0.0	0.0
65–69	4	4	8	13.9	13.2	13.5	0	0	0	0.0	0.0	0.0
70–74	3	2	5	11.3	6.7	8.9	0	0	0	0.0	0.0	0.0
75–79	4	1	5	18.5	3.6	10.1	1	0	1	4.6	0.0	2.0
80–84	1	2	3	7.8	9.9	9.1	1	0	1	7.8	0.0	3.0
85+	3	5	8	37.4	27.4	30.4	0	1	1	0.0	5.5	3.8
Total/Crude rate	37	25	62	4.9	3.3	4.1	2	1	3	0.3	0.1	0.2
AS Rate (Aust)				4.8	2.7	3.7				0.3	0.1	0.2
Confidence Interval (95%)				3.3 – 6.4	1.6 – 3.8	2.8 – 4.6				0.0 – 0.6	0.0 – 0.2	0.0 – 0.3
AS Rate (World)				3.7	2.1	2.8				0.1	0.0	0.1
Confidence Interval (95%)				2.6 – 4.8	1.3 – 2.9	2.2 – 3.5				0.0 – 0.3	0.0 – 0.1	0.0 – 0.1
Risk				1 in 273	1 in 456	1 in 343				N/A	N/A	N/A
PYLL										0	0	0
% of all cancers	0.9	0.7	0.8				0.9	0.7	0.8	0.1	0.1	0.1
Trend 1998–2002 (per annum)				-10.7%	-8.3%	-9.9%				13.5%	undefined	30.1%

Table A59 – Cancer of the Skin of lip (173.0)

South Australia Year – 2002	Incidence						Mortality					
	New cases			Rates			Deaths			Rates		
	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons
0–4	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
5–9	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
10–14	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
15–19	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
20–24	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
25–29	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
30–34	2	0	2	3.6	0.0	1.8	0	0	0	0.0	0.0	0.0
35–39	1	1	2	1.8	1.8	1.8	0	0	0	0.0	0.0	0.0
40–44	3	0	3	5.1	0.0	2.6	0	0	0	0.0	0.0	0.0
45–49	2	1	3	3.7	1.8	2.8	0	0	0	0.0	0.0	0.0
50–54	4	1	5	7.8	1.9	4.8	0	0	0	0.0	0.0	0.0
55–59	2	4	6	4.5	9.0	6.8	0	0	0	0.0	0.0	0.0
60–64	6	1	7	17.6	2.9	10.2	0	0	0	0.0	0.0	0.0
65–69	3	3	6	10.4	9.9	10.2	0	0	0	0.0	0.0	0.0
70–74	2	2	4	7.5	6.7	7.1	0	0	0	0.0	0.0	0.0
75–79	3	1	4	13.9	3.6	8.1	1	0	1	4.6	0.0	2.0
80–84	1	2	3	7.8	9.9	9.1	0	0	0	0.0	0.0	0.0
85+	2	5	7	24.9	27.4	26.6	0	1	1	0.0	5.5	3.8
Total/Crude rate	31	21	52	4.1	2.7	3.4	1	1	2	0.1	0.1	0.1
AS Rate (Aust)				4.0	2.2	3.1				0.1	0.1	0.1
Confidence Interval (95%)				2.6 – 5.4	1.2 – 3.1	2.3 – 3.9				0.0 – 0.4	0.0 – 0.2	0.0 – 0.3
AS Rate (World)				3.2	1.6	2.4				0.1	0.0	0.1
Confidence Interval (95%)				2.1 – 4.2	0.9 – 2.3	1.7 – 3.0				0.0 – 0.2	0.0 – 0.1	0.0 – 0.1
Risk				1 in 322	1 in 588	1 in 418				N/A	N/A	N/A
PYLL										0	0	0
% of all cancers	0.8	0.6	0.7				0.8	0.6	0.7	0.1	0.1	0.1
Trend 1998–2002 (per annum)				-12.2%	-10.1%	-11.5%				20.4%	undefined	79.1%

Table A60 – Cancer of the Skin of anus (173.5)

South Australia Year – 2002	Incidence						Mortality					
	New cases			Rates			Deaths			Rates		
	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons
0–4	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
5–9	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
10–14	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
15–19	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
20–24	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
25–29	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
30–34	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
35–39	0	1	1	0.0	1.8	0.9	0	0	0	0.0	0.0	0.0
40–44	1	0	1	1.7	0.0	0.9	0	0	0	0.0	0.0	0.0
45–49	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
50–54	0	1	1	0.0	1.9	1.0	0	0	0	0.0	0.0	0.0
55–59	1	0	1	2.3	0.0	1.1	0	0	0	0.0	0.0	0.0
60–64	0	1	1	0.0	2.9	1.5	0	0	0	0.0	0.0	0.0
65–69	1	1	2	3.5	3.3	3.4	0	0	0	0.0	0.0	0.0
70–74	1	0	1	3.8	0.0	1.8	0	0	0	0.0	0.0	0.0
75–79	1	0	1	4.6	0.0	2.0	0	0	0	0.0	0.0	0.0
80–84	0	0	0	0.0	0.0	0.0	1	0	1	7.8	0.0	3.0
85+	1	0	1	12.5	0.0	3.8	0	0	0	0.0	0.0	0.0
Total/Crude rate	6	4	10	0.8	0.5	0.7	1	0	1	0.1	0.0	0.1
AS Rate (Aust)				0.8	0.5	0.6				0.1	0.0	0.1
Confidence Interval (95%)				0.2 – 1.4	0.0 – 1.0	0.2 – 1.0				0.0 – 0.4	0.0 – 0.0	0.0 – 0.2
AS Rate (World)				0.5	0.4	0.5				0.1	0.0	0.0
Confidence Interval (95%)				0.2 – 0.9	0.0 – 0.8	0.2 – 0.8				0.0 – 0.1	0.0 – 0.0	0.0 – 0.1
Risk				1 in 1,783	1 in 2,025	1 in 1,914				N/A	N/A	N/A
PYLL										0	0	0
% of all cancers	0.1	0.1	0.1				0.1	0.1	0.1	0.1	0.0	0.0
Trend 1998–2002 (per annum)				18.4%	24.6%	18.6%				5.4%	0.0%	-5.5%

Note: All rates are expressed per 100,000 population. Age-standardised rates (AS Rate) use the Australian 2001 and New World Standard populations (see Methods)

Table A61 – Cancer of the Female breast (174)

South Australia Year – 2002	Incidence						Mortality					
	New cases			Rates			Deaths			Rates		
	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons
0-4	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
5-9	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
10-14	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
15-19	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
20-24	0	1	1	0.0	2.1	1.0	0	0	0	0.0	0.0	0.0
25-29	0	4	4	0.0	8.4	4.1	0	0	0	0.0	0.0	0.0
30-34	0	15	15	0.0	27.8	13.7	0	0	0	0.0	0.0	0.0
35-39	0	29	29	0.0	52.3	26.0	0	11	11	0.0	19.8	9.9
40-44	0	70	70	0.0	118.9	59.7	0	8	8	0.0	13.6	6.8
45-49	0	87	87	0.0	159.0	80.3	0	17	17	0.0	31.1	15.7
50-54	0	152	152	0.0	288.0	145.6	0	34	34	0.0	64.4	32.6
55-59	0	141	141	0.0	317.4	159.3	0	31	31	0.0	69.8	35.0
60-64	0	136	136	0.0	391.8	197.7	0	25	25	0.0	72.0	36.3
65-69	0	131	131	0.0	432.5	221.6	0	32	32	0.0	105.7	54.1
70-74	0	77	77	0.0	259.0	136.8	0	16	16	0.0	53.8	28.4
75-79	0	91	91	0.0	326.8	184.1	0	30	30	0.0	107.8	60.7
80-84	0	57	57	0.0	283.2	172.9	0	36	36	0.0	178.9	109.2
85+	0	49	49	0.0	268.1	186.3	0	50	50	0.0	273.5	190.1
Total/Crude rate	0	1,040	1,040	0.0	135.5	68.5	0	290	290	0.0	37.8	19.1
AS Rate (Aust)				0.0	120.4	62.7				0.0	30.9	16.8
Confidence Interval (95%)				0.0 – 0.0	113.0 – 127.8	58.9 – 66.5				0.0 – 0.0	27.3 – 34.5	14.9 – 18.8
AS Rate (World)				0.0	96.5	49.7				0.0	22.8	12.1
Confidence Interval (95%)				0.0 – 0.0	90.7 – 102.2	46.8 – 52.6				0.0 – 0.0	20.1 – 25.4	10.8 – 13.4
Risk				N/A	1 in 10	1 in 20				N/A	1 in 47	1 in 92
PYLL										0	3,040	3,040
% of all cancers	0.0	28.8	13.5				0.0	28.8	13.5	0.0	19.3	8.7
Trend 1998-2002 (per annum)				0.0%	-0.1%	-0.3%				0.0%	2.2%	2.1%

Table A62 – Cancer of the Male breast (175)

South Australia Year – 2002	Incidence						Mortality					
	New cases			Rates			Deaths			Rates		
	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons
0-4	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
5-9	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
10-14	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
15-19	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
20-24	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
25-29	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
30-34	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
35-39	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
40-44	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
45-49	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
50-54	1	0	1	1.9	0.0	1.0	0	0	0	0.0	0.0	0.0
55-59	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
60-64	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
65-69	3	0	3	10.4	0.0	5.1	0	0	0	0.0	0.0	0.0
70-74	1	0	1	3.8	0.0	1.8	1	0	1	3.8	0.0	1.8
75-79	1	0	1	4.6	0.0	2.0	0	0	0	0.0	0.0	0.0
80-84	1	0	1	7.8	0.0	3.0	0	0	0	0.0	0.0	0.0
85+	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
Total/Crude rate	7	0	7	0.9	0.0	0.5	1	0	1	0.1	0.0	0.1
AS Rate (Aust)				0.9	0.0	0.4				0.1	0.0	0.1
Confidence Interval (95%)				0.2 – 1.5	0.0 – 0.0	0.1 – 0.7				0.0 – 0.4	0.0 – 0.0	0.0 – 0.2
AS Rate (World)				0.6	0.0	0.3				0.1	0.0	0.0
Confidence Interval (95%)				0.2 – 1.1	0.0 – 0.0	0.1 – 0.5				0.0 – 0.2	0.0 – 0.0	0.0 – 0.1
Risk				1 in 1,242	N/A	1 in 2,561				1 in 5,312	N/A	1 in 11,257
PYLL										3	0	3
% of all cancers	0.2	0.0	0.1				0.2	0.0	0.1	0.1	0.0	0.0
Trend 1998-2002 (per annum)				2.9%	0.0%	3.1%				13.4%	0.0%	8.9%

Table A63 – Cancer of the Cervix uteri, unspecified (180.9)

South Australia Year – 2002	Incidence						Mortality					
	New cases			Rates			Deaths			Rates		
	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons
0-4	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
5-9	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
10-14	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
15-19	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
20-24	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
25-29	0	4	4	0.0	8.4	4.1	0	0	0	0.0	0.0	0.0
30-34	0	3	3	0.0	5.6	2.7	0	0	0	0.0	0.0	0.0
35-39	0	4	4	0.0	7.2	3.6	0	1	1	0.0	1.8	0.9
40-44	0	6	6	0.0	10.2	5.1	0	2	2	0.0	3.4	1.7
45-49	0	4	4	0.0	7.3	3.7	0	1	1	0.0	1.8	0.9
50-54	0	4	4	0.0	7.6	3.8	0	5	5	0.0	9.5	4.8
55-59	0	4	4	0.0	9.0	4.5	0	0	0	0.0	0.0	0.0
60-64	0	2	2	0.0	5.8	2.9	0	2	2	0.0	5.8	2.9
65-69	0	2	2	0.0	6.6	3.4	0	0	0	0.0	0.0	0.0
70-74	0	1	1	0.0	3.4	1.8	0	1	1	0.0	3.4	1.8
75-79	0	2	2	0.0	7.2	4.0	0	1	1	0.0	3.6	2.0
80-84	0	1	1	0.0	5.0	3.0	0	2	2	0.0	9.9	6.1
85+	0	3	3	0.0	16.4	11.4	0	4	4	0.0	21.9	15.2
Total/Crude rate	0	40	40	0.0	5.2	2.6	0	19	19	0.0	2.5	1.3
AS Rate (Aust)				0.0	4.9	2.5				0.0	2.1	1.1
Confidence Interval (95%)				0.0 – 0.0	3.4 – 6.5	1.7 – 3.3				0.0 – 0.0	1.1 – 3.0	0.6 – 1.6
AS Rate (World)				0.0	4.3	2.2				0.0	1.5	0.8
Confidence Interval (95%)				0.0 – 0.0	3.0 – 5.6	1.5 – 2.8				0.0 – 0.0	0.8 – 2.2	0.5 – 1.2
Risk				N/A	1 in 282	1 in 561				N/A	1 in 781	1 in 1,539
PYLL										0	270	270
% of all cancers	0.0	1.1	0.5				0.0	1.1	0.5	0.0	1.3	0.6
Trend 1998-2002 (per annum)				0.0%	-8.1%	-8.1%				0.0%	9.1%	9.3%

Note: All rates are expressed per 100,000 population. Age-standardised rates (AS Rate) use the Australian 2001 and New World Standard populations (see Methods)

Table A64 – Cancer of the Placenta (181)

South Australia Year – 2002	Incidence						Mortality					
	New cases			Rates			Deaths			Rates		
	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons
0–4	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
5–9	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
10–14	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
15–19	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
20–24	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
25–29	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
30–34	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
35–39	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
40–44	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
45–49	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
50–54	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
55–59	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
60–64	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
65–69	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
70–74	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
75–79	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
80–84	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
85+	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
Total/Crude rate	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
AS Rate (Aust)				0.0	0.0	0.0				0.0	0.0	0.0
Confidence Interval (95%)				0.0 – 0.0	0.0 – 0.0	0.0 – 0.0				0.0 – 0.0	0.0 – 0.0	0.0 – 0.0
AS Rate (World)				0.0	0.0	0.0				0.0	0.0	0.0
Confidence Interval (95%)				0.0 – 0.0	0.0 – 0.0	0.0 – 0.0				0.0 – 0.0	0.0 – 0.0	0.0 – 0.0
Risk				N/A	N/A	N/A				N/A	N/A	N/A
PYLL										0	0	0
% of all cancers	0.0	0.0	0.0				0.0	0.0	0.0	0.0	0.0	0.0
Trend 1998–2002 (per annum)				0.0%	-7.5%	-7.8%				0.0%	0.0%	0.0%

Table A65 – Cancer of the Body of the uterus (182)

South Australia Year – 2002	Incidence						Mortality					
	New cases			Rates			Deaths			Rates		
	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons
0–4	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
5–9	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
10–14	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
15–19	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
20–24	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
25–29	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
30–34	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
35–39	0	1	1	0.0	1.8	0.9	0	0	0	0.0	0.0	0.0
40–44	0	2	2	0.0	3.4	1.7	0	0	0	0.0	0.0	0.0
45–49	0	9	9	0.0	16.4	8.3	0	1	1	0.0	1.8	0.9
50–54	0	15	15	0.0	28.4	14.4	0	0	0	0.0	0.0	0.0
55–59	0	21	21	0.0	47.3	23.7	0	2	2	0.0	4.5	2.3
60–64	0	26	26	0.0	74.9	37.8	0	3	3	0.0	8.6	4.4
65–69	0	20	20	0.0	66.0	33.8	0	3	3	0.0	9.9	5.1
70–74	0	26	26	0.0	87.5	46.2	0	4	4	0.0	13.5	7.1
75–79	0	9	9	0.0	32.3	18.2	0	2	2	0.0	7.2	4.0
80–84	0	10	10	0.0	49.7	30.3	0	3	3	0.0	14.9	9.1
85+	0	13	13	0.0	71.1	49.4	0	8	8	0.0	43.8	30.4
Total/Crude rate	0	152	152	0.0	17.0	9.0	0	26	26	0.0	3.4	1.7
AS Rate (Aust)				0.0	17.0	9.0				0.0	2.6	1.5
Confidence Interval (95%)				0.0 – 0.0	14.2 – 19.7	7.5 – 10.4				0.0 – 0.0	1.6 – 3.6	0.9 – 2.0
AS Rate (World)				0.0	13.1	6.8				0.0	1.7	1.0
Confidence Interval (95%)				0.0 – 0.0	11.0 – 15.2	5.7 – 7.9				0.0 – 0.0	1.0 – 2.4	0.6 – 1.3
Risk				N/A	1 in 62	1 in 120				N/A	1 in 522	1 in 1,014
PYLL										0	133	133
% of all cancers	0.0	4.2	2.0				0.0	4.2	2.0	0.0	1.7	0.8
Trend 1998–2002 (per annum)				0.0%	-1.5%	-1.6%				0.0%	-1.7%	-2.1%

Table A66 – Cancer of the Ovary and other uterine adnexa (183)

South Australia Year – 2002	Incidence						Mortality					
	New cases			Rates			Deaths			Rates		
	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons
0–4	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
5–9	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
10–14	0	1	1	0.0	2.0	1.0	0	0	0	0.0	0.0	0.0
15–19	0	1	1	0.0	2.0	1.0	0	0	0	0.0	0.0	0.0
20–24	0	1	1	0.0	2.1	1.0	0	0	0	0.0	0.0	0.0
25–29	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
30–34	0	2	2	0.0	3.7	1.8	0	1	1	0.0	1.9	0.9
35–39	0	2	2	0.0	3.6	1.8	0	0	0	0.0	0.0	0.0
40–44	0	4	4	0.0	6.8	3.4	0	1	1	0.0	1.7	0.9
45–49	0	6	6	0.0	11.0	5.5	0	0	0	0.0	0.0	0.0
50–54	0	6	6	0.0	11.4	5.7	0	8	8	0.0	15.2	7.7
55–59	0	10	10	0.0	22.5	11.3	0	6	6	0.0	13.5	6.8
60–64	0	13	13	0.0	37.4	18.9	0	12	12	0.0	34.6	17.4
65–69	0	11	11	0.0	36.3	18.6	0	12	12	0.0	39.6	20.3
70–74	0	9	9	0.0	30.3	16.0	0	8	8	0.0	26.9	14.2
75–79	0	16	16	0.0	57.5	32.4	0	11	11	0.0	39.5	22.3
80–84	0	11	11	0.0	54.7	33.4	0	9	9	0.0	44.7	27.3
85+	0	11	11	0.0	60.2	41.8	0	3	3	0.0	16.4	11.4
Total/Crude rate	0	104	104	0.0	13.6	6.8	0	71	71	0.0	9.3	4.7
AS Rate (Aust)				0.0	11.3	6.1				0.0	7.8	4.1
Confidence Interval (95%)				0.0 – 0.0	9.1 – 13.6	4.9 – 7.3				0.0 – 0.0	5.9 – 9.6	3.2 – 5.1
AS Rate (World)				0.0	8.7	4.6				0.0	5.8	3.1
Confidence Interval (95%)				0.0 – 0.0	7.0 – 10.4	3.7 – 5.4				0.0 – 0.0	4.5 – 7.2	2.4 – 3.8
Risk				N/A	1 in 119	1 in 233				N/A	1 in 151	1 in 294
PYLL										0	620	620
% of all cancers	0.0	2.9	1.3				0.0	2.9	1.3	0.0	4.7	2.1
Trend 1998–2002 (per annum)				0.0%	-2.7%	-2.5%				0.0%	2.2%	1.7%

Note: All rates are expressed per 100,000 population. Age-standardised rates (AS Rate) use the Australian 2001 and New World Standard populations (see Methods)

Table A67 – Cancer of other & unspecified female genitalia (184)

South Australia Year – 2002	Incidence						Mortality					
	New cases			Rates			Deaths			Rates		
	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons
0-4	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
5-9	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
10-14	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
15-19	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
20-24	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
25-29	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
30-34	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
35-39	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
40-44	0	1	1	0.0	1.7	0.9	0	0	0	0.0	0.0	0.0
45-49	0	1	1	0.0	1.8	0.9	0	0	0	0.0	0.0	0.0
50-54	0	0	0	0.0	0.0	0.0	0	1	1	0.0	1.9	1.0
55-59	0	1	1	0.0	2.3	1.1	0	1	1	0.0	2.3	1.1
60-64	0	3	3	0.0	8.6	4.4	0	1	1	0.0	2.9	1.5
65-69	0	1	1	0.0	3.3	1.7	0	0	0	0.0	0.0	0.0
70-74	0	2	2	0.0	6.7	3.6	0	0	0	0.0	0.0	0.0
75-79	0	3	3	0.0	10.8	6.1	0	2	2	0.0	7.2	4.0
80-84	0	2	2	0.0	9.9	6.1	0	3	3	0.0	14.9	9.1
85+	0	4	4	0.0	21.9	15.2	0	5	5	0.0	27.4	19.0
Total/Crude rate	0	18	18	0.0	2.3	1.2	0	13	13	0.0	1.7	0.9
AS Rate (Aust)				0.0	1.8	1.0				0.0	1.2	0.7
Confidence Interval (95%)				0.0 – 0.0	1.0 – 2.7	0.5 – 1.5				0.0 – 0.0	0.5 – 1.8	0.3 – 1.1
AS Rate (World)				0.0	1.3	0.7				0.0	0.7	0.4
Confidence Interval (95%)				0.0 – 0.0	0.7 – 1.9	0.4 – 1.0				0.0 – 0.0	0.3 – 1.1	0.2 – 0.6
Risk				N/A	1 in 819	1 in 1,599				N/A	1 in 2,847	1 in 5,648
PYLL										0	53	53
% of all cancers	0.0	0.5	0.2				0.0	0.5	0.2	0.0	0.9	0.4
Trend 1998–2002 (per annum)				0.0%	-3.8%	-4.4%				0.0%	-0.3%	-1.5%

Table A68 – Cancer of the Vagina & unspec female genital organs (184.0 184.8 184.9)

South Australia Year – 2002	Incidence						Mortality					
	New cases			Rates			Deaths			Rates		
	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons
0-4	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
5-9	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
10-14	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
15-19	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
20-24	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
25-29	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
30-34	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
35-39	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
40-44	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
45-49	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
50-54	0	0	0	0.0	0.0	0.0	0	1	1	0.0	1.9	1.0
55-59	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
60-64	0	0	0	0.0	0.0	0.0	0	1	1	0.0	2.9	1.5
65-69	0	1	1	0.0	3.3	1.7	0	0	0	0.0	0.0	0.0
70-74	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
75-79	0	1	1	0.0	3.6	2.0	0	1	1	0.0	3.6	2.0
80-84	0	1	1	0.0	5.0	3.0	0	0	0	0.0	0.0	0.0
85+	0	0	0	0.0	0.0	0.0	0	1	1	0.0	5.5	3.8
Total/Crude rate	0	3	3	0.0	0.4	0.2	0	4	4	0.0	0.5	0.3
AS Rate (Aust)				0.0	0.3	0.2				0.0	0.4	0.2
Confidence Interval (95%)				0.0 – 0.0	0.0 – 0.6	0.0 – 0.4				0.0 – 0.0	0.0 – 0.8	0.0 – 0.5
AS Rate (World)				0.0	0.2	0.1				0.0	0.3	0.2
Confidence Interval (95%)				0.0 – 0.0	0.0 – 0.4	0.0 – 0.2				0.0 – 0.0	0.0 – 0.6	0.0 – 0.3
Risk				N/A	1 in 6,058	1 in 11,821				N/A	1 in 4,189	1 in 8,294
PYLL										0	35	35
% of all cancers	0.0	0.1	0.0				0.0	0.1	0.0	0.0	0.3	0.1
Trend 1998–2002 (per annum)				0.0%	-4.3%	-3.9%				0.0%	2.0%	0.7%

Table A69 – Cancer of the Vulva (184.1 184.2 184.3 184.4)

South Australia Year – 2002	Incidence						Mortality					
	New cases			Rates			Deaths			Rates		
	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons
0-4	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
5-9	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
10-14	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
15-19	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
20-24	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
25-29	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
30-34	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
35-39	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
40-44	0	1	1	0.0	1.7	0.9	0	0	0	0.0	0.0	0.0
45-49	0	1	1	0.0	1.8	0.9	0	0	0	0.0	0.0	0.0
50-54	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
55-59	0	1	1	0.0	2.3	1.1	0	1	1	0.0	2.3	1.1
60-64	0	3	3	0.0	8.6	4.4	0	0	0	0.0	0.0	0.0
65-69	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
70-74	0	2	2	0.0	6.7	3.6	0	0	0	0.0	0.0	0.0
75-79	0	2	2	0.0	7.2	4.0	0	1	1	0.0	3.6	2.0
80-84	0	1	1	0.0	5.0	3.0	0	3	3	0.0	14.9	9.1
85+	0	4	4	0.0	21.9	15.2	0	4	4	0.0	21.9	15.2
Total/Crude rate	0	15	15	0.0	2.0	1.0	0	9	9	0.0	1.2	0.6
AS Rate (Aust)				0.0	1.5	0.9				0.0	0.8	0.5
Confidence Interval (95%)				0.0 – 0.0	0.7 – 2.3	0.4 – 1.3				0.0 – 0.0	0.3 – 1.3	0.2 – 0.8
AS Rate (World)				0.0	1.1	0.6				0.0	0.4	0.3
Confidence Interval (95%)				0.0 – 0.0	0.5 – 1.7	0.3 – 0.9				0.0 – 0.0	0.2 – 0.7	0.1 – 0.4
Risk				N/A	1 in 946	1 in 1,849				N/A	1 in 8,885	1 in 17,705
PYLL										0	18	18
% of all cancers	0.0	0.4	0.2				0.0	0.4	0.2	0.0	0.6	0.3
Trend 1998–2002 (per annum)				0.0%	-3.7%	-4.6%				0.0%	-1.9%	-2.8%

Note: All rates are expressed per 100,000 population. Age-standardised rates (AS Rate) use the Australian 2001 and New World Standard populations (see Methods)

Table A70 – Cancer of the Prostate (185)

South Australia Year – 2002	Incidence						Mortality					
	New cases			Rates			Deaths			Rates		
	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons
0–4	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
5–9	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
10–14	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
15–19	1	0	1	1.9	0.0	1.0	0	0	0	0.0	0.0	0.0
20–24	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
25–29	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
30–34	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
35–39	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
40–44	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
45–49	13	0	13	24.2	0.0	12.0	1	0	1	1.9	0.0	0.9
50–54	30	0	30	58.1	0.0	28.7	0	0	0	0.0	0.0	0.0
55–59	79	0	79	179.1	0.0	89.2	5	0	5	11.3	0.0	5.6
60–64	121	0	121	355.0	0.0	175.9	9	0	9	26.4	0.0	13.1
65–69	163	0	163	565.7	0.0	275.8	10	0	10	34.7	0.0	16.9
70–74	196	0	196	738.0	0.0	348.3	29	0	29	109.2	0.0	51.5
75–79	158	0	158	732.0	0.0	319.7	47	0	47	217.8	0.0	95.1
80–84	99	0	99	770.4	0.0	300.2	48	0	48	373.5	0.0	145.6
85+	86	0	86	1,071.9	0.0	327.0	73	0	73	909.9	0.0	277.5
Total/Crude rate	946	0	946	125.9	0.0	62.3	222	0	222	29.5	0.0	14.6
AS Rate (Aust)				121.5	0.0	54.2				31.3	0.0	12.0
Confidence Interval (95%)				113.8 – 129.3	0.0 – 0.0	50.7 – 57.6				27.1 – 35.4	0.0 – 0.0	10.4 – 13.6
AS Rate (World)				83.7	0.0	38.4				17.1	0.0	6.8
Confidence Interval (95%)				78.7 – 88.7	0.0 – 0.0	36.0 – 40.7				15.3 – 19.0	0.0 – 0.0	6.1 – 7.6
Risk				1 in 11	N/A	1 in 22				1 in 109	N/A	1 in 228
PYLL										375	0	375
% of all cancers	23.1	0.0	12.3				23.1	0.0	12.3	12.2	0.0	6.7
Trend 1998–2002 (per annum)				-2.6%	0.0%	-2.1%				-1.9%	0.0%	-2.5%

Table A71 – Cancer of the Testis (186)

South Australia Year – 2002	Incidence						Mortality					
	New cases			Rates			Deaths			Rates		
	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons
0–4	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
5–9	1	0	1	2.0	0.0	1.0	0	0	0	0.0	0.0	0.0
10–14	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
15–19	4	0	4	7.6	0.0	3.9	0	0	0	0.0	0.0	0.0
20–24	5	0	5	10.1	0.0	5.2	0	0	0	0.0	0.0	0.0
25–29	6	0	6	12.1	0.0	6.2	0	0	0	0.0	0.0	0.0
30–34	9	0	9	16.3	0.0	8.2	0	0	0	0.0	0.0	0.0
35–39	8	0	8	14.3	0.0	7.2	0	0	0	0.0	0.0	0.0
40–44	4	0	4	6.9	0.0	3.4	0	0	0	0.0	0.0	0.0
45–49	2	0	2	3.7	0.0	1.8	0	0	0	0.0	0.0	0.0
50–54	1	0	1	1.9	0.0	1.0	0	0	0	0.0	0.0	0.0
55–59	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
60–64	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
65–69	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
70–74	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
75–79	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
80–84	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
85+	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
Total/Crude rate	40	0	40	5.3	0.0	2.6	0	0	0	0.0	0.0	0.0
AS Rate (Aust)				5.5	0.0	2.8				0.0	0.0	0.0
Confidence Interval (95%)				3.8 – 7.2	0.0 – 0.0	1.9 – 3.6				0.0 – 0.0	0.0 – 0.0	0.0 – 0.0
AS Rate (World)				5.6	0.0	2.9				0.0	0.0	0.0
Confidence Interval (95%)				4.0 – 7.3	0.0 – 0.0	2.0 – 3.7				0.0 – 0.0	0.0 – 0.0	0.0 – 0.0
Risk				1 in 268	N/A	1 in 528				N/A	N/A	N/A
PYLL										0	0	0
% of all cancers	1.0	0.0	0.5				1.0	0.0	0.5	0.0	0.0	0.0
Trend 1998–2002 (per annum)				-1.7%	0.0%	-1.5%				-17.4%	0.0%	-15.7%

Table A72 – Cancer of the Penis and other male genital organs (187)

South Australia Year – 2002	Incidence						Mortality					
	New cases			Rates			Deaths			Rates		
	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons
0–4	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
5–9	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
10–14	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
15–19	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
20–24	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
25–29	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
30–34	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
35–39	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
40–44	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
45–49	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
50–54	2	0	2	3.9	0.0	1.9	0	0	0	0.0	0.0	0.0
55–59	0	0	0	0.0	0.0	0.0	1	0	1	2.3	0.0	1.1
60–64	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
65–69	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
70–74	3	0	3	11.3	0.0	5.3	0	0	0	0.0	0.0	0.0
75–79	1	0	1	4.6	0.0	2.0	0	0	0	0.0	0.0	0.0
80–84	1	0	1	7.8	0.0	3.0	0	0	0	0.0	0.0	0.0
85+	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
Total/Crude rate	7	0	7	0.9	0.0	0.5	1	0	1	0.1	0.0	0.1
AS Rate (Aust)				0.9	0.0	0.4				0.1	0.0	0.1
Confidence Interval (95%)				0.2 – 1.5	0.0 – 0.0	0.1 – 0.7				0.0 – 0.3	0.0 – 0.0	0.0 – 0.2
AS Rate (World)				0.6	0.0	0.3				0.1	0.0	0.1
Confidence Interval (95%)				0.2 – 1.0	0.0 – 0.0	0.1 – 0.5				0.0 – 0.3	0.0 – 0.0	0.0 – 0.1
Risk				1 in 1,319	N/A	1 in 2,760				1 in 8,821	N/A	1 in 17,705
PYLL										18	0	18
% of all cancers	0.2	0.0	0.1				0.2	0.0	0.1	0.1	0.0	0.0
Trend 1998–2002 (per annum)				-8.5%	0.0%	-7.0%				78.9%	0.0%	116.3%

Note: All rates are expressed per 100,000 population. Age-standardised rates (AS Rate) use the Australian 2001 and New World Standard populations (see Methods)

Table A73 – Cancer of the Bladder (188)

South Australia Year – 2002	Incidence						Mortality					
	New cases			Rates			Deaths			Rates		
	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons
0–4	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
5–9	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
10–14	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
15–19	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
20–24	1	0	1	2.0	0.0	1.0	0	0	0	0.0	0.0	0.0
25–29	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
30–34	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
35–39	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
40–44	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
45–49	2	1	3	3.7	1.8	2.8	1	0	1	1.9	0.0	0.9
50–54	5	3	8	9.7	5.7	7.7	4	0	4	7.8	0.0	3.8
55–59	8	1	9	18.1	2.3	10.2	1	1	2	2.3	2.3	2.3
60–64	9	4	13	26.4	11.5	18.9	1	0	1	2.9	0.0	1.5
65–69	15	2	17	52.1	6.6	28.8	7	1	8	24.3	3.3	13.5
70–74	24	3	27	90.4	10.1	48.0	6	3	9	22.6	10.1	16.0
75–79	18	7	25	83.4	25.1	50.6	13	7	20	60.2	25.1	40.5
80–84	29	14	43	225.7	69.6	130.4	17	6	23	132.3	29.8	69.7
85+	13	11	24	162.0	60.2	91.2	14	1	15	174.5	5.5	57.0
Total/Crude rate	124	46	170	16.5	6.0	11.2	64	19	83	8.5	2.5	5.5
AS Rate (Aust)				16.2	4.4	9.5				8.7	1.8	4.5
Confidence Interval (95%)				13.3 – 19.1	3.1 – 5.6	8.1 – 10.9				6.6 – 10.9	1.0 – 2.6	3.6 – 5.5
AS Rate (World)				10.5	2.7	6.2				5.1	1.1	2.8
Confidence Interval (95%)				8.9 – 12.2	1.9 – 3.5	5.3 – 7.1				4.1 – 6.2	0.7 – 1.5	2.2 – 3.3
Risk				1 in 99	1 in 527	1 in 171				1 in 325	1 in 1,279	1 in 527
PYLL										215	33	248
% of all cancers	3.0	1.3	2.2				3.0	1.3	2.2	3.5	1.3	2.5
Trend 1998–2002 (per annum)				-2.4%	-1.1%	-1.8%				3.1%	-7.2%	0.0%

Table A74 – Cancer of the Kidney, and other unspec urinary organs (189)

South Australia Year – 2002	Incidence						Mortality					
	New cases			Rates			Deaths			Rates		
	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons
0–4	0	1	1	0.0	2.3	1.1	0	0	0	0.0	0.0	0.0
5–9	0	0	0	0.0	0.0	0.0	0	1	1	0.0	2.1	1.0
10–14	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
15–19	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
20–24	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
25–29	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
30–34	1	0	1	1.8	0.0	0.9	0	0	0	0.0	0.0	0.0
35–39	3	3	6	5.4	5.4	5.4	0	1	1	0.0	1.8	0.9
40–44	3	3	6	5.1	5.1	5.1	0	1	1	0.0	1.7	0.9
45–49	7	4	11	13.0	7.3	10.1	1	0	1	1.9	0.0	0.9
50–54	9	9	18	17.4	17.1	17.2	3	3	6	5.8	5.7	5.7
55–59	14	6	20	31.7	13.5	22.6	6	2	8	13.6	4.5	9.0
60–64	15	5	20	44.0	14.4	29.1	7	2	9	20.5	5.8	13.1
65–69	15	2	17	52.1	6.6	28.8	7	1	8	24.3	3.3	13.5
70–74	26	7	33	97.9	23.5	58.6	6	3	9	22.6	10.1	16.0
75–79	17	7	24	78.8	25.1	48.6	5	6	11	23.2	21.6	22.3
80–84	14	11	25	108.9	54.7	75.8	9	6	15	70.0	29.8	45.5
85+	7	9	16	87.2	49.2	60.8	6	8	14	74.8	43.8	53.2
Total/Crude rate	131	67	198	17.4	8.7	13.0	50	34	84	6.7	4.4	5.5
AS Rate (Aust)				16.7	7.2	11.6				6.5	3.4	4.8
Confidence Interval (95%)				13.9 – 19.6	5.4 – 9.0	10.0 – 13.2				4.7 – 8.3	2.2 – 4.6	3.8 – 5.8
AS Rate (World)				12.1	5.3	8.5				4.5	2.3	3.3
Confidence Interval (95%)				10.2 – 14.0	4.0 – 6.6	7.3 – 9.6				3.3 – 5.6	1.5 – 3.1	2.6 – 4.0
Risk				1 in 75	1 in 211	1 in 112				1 in 226	1 in 573	1 in 328
PYLL										355	280	635
% of all cancers	3.2	1.9	2.6				3.2	1.9	2.6	2.7	2.3	2.5
Trend 1998–2002 (per annum)				0.7%	-0.7%	0.3%				2.2%	1.9%	1.9%

Table A75 – Cancer of the Eye (190)

South Australia Year – 2002	Incidence						Mortality					
	New cases			Rates			Deaths			Rates		
	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons
0–4	0	2	2	0.0	4.5	2.2	0	0	0	0.0	0.0	0.0
5–9	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
10–14	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
15–19	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
20–24	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
25–29	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
30–34	0	2	2	0.0	3.7	1.8	0	0	0	0.0	0.0	0.0
35–39	0	1	1	0.0	1.8	0.9	0	0	0	0.0	0.0	0.0
40–44	1	0	1	1.7	0.0	0.9	0	0	0	0.0	0.0	0.0
45–49	1	1	2	1.9	1.8	1.8	2	0	2	3.7	0.0	1.8
50–54	0	2	2	0.0	3.8	1.9	1	0	1	1.9	0.0	1.0
55–59	9	0	9	20.4	0.0	10.2	0	0	0	0.0	0.0	0.0
60–64	1	0	1	2.9	0.0	1.5	0	0	0	0.0	0.0	0.0
65–69	1	2	3	3.5	6.6	5.1	0	0	0	0.0	0.0	0.0
70–74	3	2	5	11.3	6.7	8.9	0	0	0	0.0	0.0	0.0
75–79	2	0	2	9.3	0.0	4.0	1	1	2	4.6	3.6	4.0
80–84	2	0	2	15.6	0.0	6.1	0	0	0	0.0	0.0	0.0
85+	1	2	3	12.5	10.9	11.4	0	0	0	0.0	0.0	0.0
Total/Crude rate	21	14	35	2.8	1.8	2.3	4	1	5	0.5	0.1	0.3
AS Rate (Aust)				2.6	1.7	2.1				0.5	0.1	0.3
Confidence Interval (95%)				1.5 – 3.7	0.8 – 2.6	1.4 – 2.8				0.0 – 1.0	0.0 – 0.3	0.0 – 0.6
AS Rate (World)				2.0	1.5	1.7				0.4	0.1	0.2
Confidence Interval (95%)				1.2 – 2.7	0.7 – 2.4	1.2 – 2.3				0.0 – 0.8	0.0 – 0.1	0.0 – 0.4
Risk				1 in 480	1 in 691	1 in 570				1 in 3,531	N/A	1 in 7,135
PYLL										78	0	78
% of all cancers	0.5	0.4	0.5				0.5	0.4	0.5	0.2	0.1	0.2
Trend 1998–2002 (per annum)				-1.2%	-0.9%	-1.3%				3.0%	-23.4%	-11.6%

Note: All rates are expressed per 100,000 population. Age-standardised rates (AS Rate) use the Australian 2001 and New World Standard populations (see Methods)

Table A76 – Cancer of the Brain (191)

South Australia Year – 2002	Incidence						Mortality					
	New cases			Rates			Deaths			Rates		
	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons
0-4	0	2	2	0.0	4.5	2.2	0	0	0	0.0	0.0	0.0
5-9	2	1	3	4.0	2.1	3.1	1	0	1	2.0	0.0	1.0
10-14	1	0	1	1.9	0.0	1.0	1	0	1	1.9	0.0	1.0
15-19	2	2	4	3.8	4.0	3.9	2	1	3	3.8	2.0	2.9
20-24	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
25-29	0	3	3	0.0	6.3	3.1	1	0	1	2.0	0.0	1.0
30-34	1	3	4	1.8	5.6	3.7	2	1	3	3.6	1.9	2.7
35-39	2	1	3	3.6	1.8	2.7	1	2	3	1.8	3.6	2.7
40-44	4	1	5	6.9	1.7	4.3	1	2	3	1.7	3.4	2.6
45-49	4	4	8	7.5	7.3	7.4	1	2	3	1.9	3.7	2.8
50-54	6	5	11	11.6	9.5	10.5	8	7	15	15.5	13.3	14.4
55-59	11	5	16	24.9	11.3	18.1	8	5	13	18.1	11.3	14.7
60-64	6	8	14	17.6	23.0	20.4	5	5	10	14.7	14.4	14.5
65-69	4	10	14	13.9	33.0	23.7	5	6	11	17.4	19.8	18.6
70-74	10	5	15	37.7	16.8	26.7	7	3	10	26.4	10.1	17.8
75-79	5	8	13	23.2	28.7	26.3	6	9	15	27.8	32.3	30.3
80-84	3	4	7	23.3	19.9	21.2	5	3	8	38.9	14.9	24.3
85+	5	3	8	62.3	16.4	30.4	4	5	9	49.9	27.4	34.2
Total/Crude rate	66	65	131	8.8	8.5	8.6	58	51	109	7.7	6.6	7.2
AS Rate (Aust)				8.5	7.6	7.9				7.5	5.7	6.5
Confidence Interval (95%)				6.5 – 10.6	5.7 – 9.5	6.6 – 9.3				5.6 – 9.5	4.1 – 7.3	5.3 – 7.7
AS Rate (World)				6.7	6.5	6.5				5.8	4.4	5.1
Confidence Interval (95%)				5.1 – 8.3	4.8 – 8.1	5.4 – 7.7				4.4 – 7.3	3.2 – 5.6	4.1 – 6.0
Risk				1 in 149	1 in 158	1 in 154				1 in 181	1 in 241	1 in 207
PYLL										913	655	1,568
% of all cancers	1.6	1.8	1.7				1.6	1.8	1.7	3.2	3.4	3.3
Trend 1998-2002 (per annum)				0.0%	7.0%	2.1%				3.0%	5.6%	3.6%

Table A77 – Cancer of the Cranial nerves, spinal cord and meninges (192)

South Australia Year – 2002	Incidence						Mortality					
	New cases			Rates			Deaths			Rates		
	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons
0-4	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
5-9	1	0	1	2.0	0.0	1.0	0	0	0	0.0	0.0	0.0
10-14	1	0	1	1.9	0.0	1.0	0	0	0	0.0	0.0	0.0
15-19	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
20-24	1	0	1	2.0	0.0	1.0	0	0	0	0.0	0.0	0.0
25-29	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
30-34	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
35-39	1	0	1	1.8	0.0	0.9	0	0	0	0.0	0.0	0.0
40-44	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
45-49	0	0	0	0.0	0.0	0.0	0	1	1	0.0	1.8	0.9
50-54	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
55-59	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
60-64	0	0	0	0.0	0.0	0.0	1	1	2	2.9	2.9	2.9
65-69	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
70-74	0	1	1	0.0	3.4	1.8	0	0	0	0.0	0.0	0.0
75-79	0	0	0	0.0	0.0	0.0	1	0	1	4.6	0.0	2.0
80-84	0	0	0	0.0	0.0	0.0	0	1	1	0.0	5.0	3.0
85+	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
Total/Crude rate	4	1	5	0.5	0.1	0.3	2	3	5	0.3	0.4	0.3
AS Rate (Aust)				0.5	0.1	0.3				0.2	0.3	0.3
Confidence Interval (95%)				0.0 – 1.1	0.0 – 0.3	0.0 – 0.6				0.0 – 0.6	0.0 – 0.7	0.0 – 0.6
AS Rate (World)				0.6	0.1	0.4				0.2	0.3	0.2
Confidence Interval (95%)				0.0 – 1.3	0.0 – 0.2	0.0 – 0.7				0.0 – 0.4	0.0 – 0.6	0.0 – 0.4
Risk				1 in 2,593	1 in 5,945	1 in 3,498				1 in 6,817	1 in 4,248	1 in 5,223
PYLL										13	40	53
% of all cancers	0.1	0.0	0.1				0.1	0.0	0.1	0.1	0.2	0.2
Trend 1998-2002 (per annum)				5.9%	-17.9%	-7.2%				-4.6%	1045.0%	36.6%

Table A78 – Cancer of the Thyroid gland (193)

South Australia Year – 2002	Incidence						Mortality					
	New cases			Rates			Deaths			Rates		
	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons
0-4	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
5-9	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
10-14	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
15-19	0	1	1	0.0	2.0	1.0	0	0	0	0.0	0.0	0.0
20-24	0	1	1	0.0	2.1	1.0	0	0	0	0.0	0.0	0.0
25-29	2	4	6	4.0	8.4	6.2	0	0	0	0.0	0.0	0.0
30-34	0	10	10	0.0	18.5	9.2	0	0	0	0.0	0.0	0.0
35-39	3	4	7	5.4	7.2	6.3	0	0	0	0.0	0.0	0.0
40-44	2	9	11	3.4	15.3	9.4	0	0	0	0.0	0.0	0.0
45-49	0	6	6	0.0	11.0	5.5	0	0	0	0.0	0.0	0.0
50-54	1	7	8	1.9	13.3	7.7	0	0	0	0.0	0.0	0.0
55-59	3	7	10	6.8	15.8	11.3	0	1	1	0.0	2.3	1.1
60-64	2	3	5	5.9	8.6	7.3	0	1	1	0.0	2.9	1.5
65-69	1	3	4	3.5	9.9	6.8	1	0	1	3.5	0.0	1.7
70-74	0	2	2	0.0	6.7	3.6	1	2	3	3.8	6.7	5.3
75-79	1	1	2	4.6	3.6	4.0	1	0	1	4.6	0.0	2.0
80-84	2	2	4	15.6	9.9	12.1	1	1	2	7.8	5.0	6.1
85+	0	2	2	0.0	10.9	7.6	2	0	2	24.9	0.0	7.6
Total/Crude rate	17	62	79	2.3	8.1	5.2	6	5	11	0.8	0.7	0.7
AS Rate (Aust)				2.2	7.8	5.0				0.8	0.5	0.6
Confidence Interval (95%)				1.2 – 3.3	5.9 – 9.8	3.9 – 6.2				0.2 – 1.5	0.1 – 1.0	0.3 – 1.0
AS Rate (World)				1.9	7.0	4.5				0.5	0.4	0.4
Confidence Interval (95%)				1.0 – 2.7	5.4 – 8.7	3.5 – 5.4				0.1 – 0.8	0.1 – 0.7	0.2 – 0.6
Risk				1 in 647	1 in 169	1 in 267				1 in 2,765	1 in 1,687	1 in 2,083
PYLL										10	35	45
% of all cancers	0.4	1.7	1.0				0.4	1.7	1.0	0.3	0.3	0.3
Trend 1998-2002 (per annum)				0.8%	6.0%	4.3%				106.1%	35.9%	41.5%

Note: All rates are expressed per 100,000 population. Age-standardised rates (AS Rate) use the Australian 2001 and New World Standard populations (see Methods)

Table A79 – Cancer of other endocrine glands (194)

South Australia Year – 2002	Incidence						Mortality					
	New cases			Rates			Deaths			Rates		
	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons
0–4	0	1	1	0.0	2.3	1.1	0	0	0	0.0	0.0	0.0
5–9	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
10–14	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
15–19	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
20–24	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
25–29	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
30–34	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
35–39	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
40–44	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
45–49	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
50–54	1	0	1	1.9	0.0	1.0	0	0	0	0.0	0.0	0.0
55–59	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
60–64	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
65–69	1	0	1	3.5	0.0	1.7	0	0	0	0.0	0.0	0.0
70–74	1	0	1	3.8	0.0	1.8	1	0	1	3.8	0.0	1.8
75–79	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
80–84	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
85+	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
Total/Crude rate	3	1	4	0.4	0.1	0.3	1	0	1	0.1	0.0	0.1
AS Rate (Aust)				0.4	0.1	0.3				0.1	0.0	0.1
Confidence Interval (95%)				0.0 – 0.8	0.0 – 0.4	0.0 – 0.5				0.0 – 0.4	0.0 – 0.0	0.0 – 0.2
AS Rate (World)				0.3	0.2	0.2				0.1	0.0	0.0
Confidence Interval (95%)				0.0 – 0.6	0.0 – 0.6	0.0 – 0.5				0.0 – 0.2	0.0 – 0.0	0.0 – 0.1
Risk				1 in 2,181	1 in 8,883	1 in 3,618				1 in 5,312	N/A	1 in 11,257
PYLL										3	0	3
% of all cancers	0.1	0.0	0.1				0.1	0.0	0.1	0.1	0.0	0.0
Trend 1998–2002 (per annum)				-7.0%	-15.1%	-12.4%				-0.3%	41.8%	14.4%

Table A80 – Cancer of Unspecified site (199)

South Australia Year – 2002	Incidence						Mortality					
	New cases			Rates			Deaths			Rates		
	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons
0–4	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
5–9	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
10–14	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
15–19	0	1	1	0.0	2.0	1.0	0	1	1	0.0	2.0	1.0
20–24	1	0	1	2.0	0.0	1.0	1	0	1	2.0	0.0	1.0
25–29	1	0	1	2.0	0.0	1.0	1	0	1	2.0	0.0	1.0
30–34	2	0	2	3.6	0.0	1.8	1	0	1	1.8	0.0	0.9
35–39	1	1	2	1.8	1.8	1.8	1	1	2	1.8	1.8	1.8
40–44	1	3	4	1.7	5.1	3.4	2	1	3	3.4	1.7	2.6
45–49	4	4	8	7.5	7.3	7.4	3	4	7	5.6	7.3	6.5
50–54	1	8	9	1.9	15.2	8.6	2	5	7	3.9	9.5	6.7
55–59	8	7	15	18.1	15.8	16.9	6	3	9	13.6	6.8	10.2
60–64	6	7	13	17.6	20.2	18.9	8	6	14	23.5	17.3	20.4
65–69	20	13	33	69.4	42.9	55.8	14	14	28	48.6	46.2	47.4
70–74	23	17	40	86.6	57.2	71.1	21	22	43	79.1	74.0	76.4
75–79	32	23	55	148.3	82.6	111.3	29	19	48	134.4	68.2	97.1
80–84	18	25	43	140.1	124.2	130.4	13	25	38	101.2	124.2	115.2
85+	8	37	45	99.7	202.4	171.1	9	32	41	112.2	175.1	155.9
Total/Crude rate	126	146	272	16.8	19.0	17.9	111	133	244	14.8	17.3	16.1
AS Rate (Aust)				16.2	14.3	15.4				14.3	13.0	13.8
Confidence Interval (95%)				13.3 – 19.0	12.0 – 16.7	13.5 – 17.2				11.6 – 17.0	10.8 – 15.3	12.0 – 15.5
AS Rate (World)				11.0	9.5	10.2				9.7	8.5	9.1
Confidence Interval (95%)				9.2 – 12.7	7.9 – 11.0	9.1 – 11.4				8.0 – 11.3	7.1 – 10.0	8.0 – 10.2
Risk				1 in 95	1 in 120	1 in 106				1 in 108	1 in 121	1 in 114
PYLL										735	638	1,373
% of all cancers	3.1	4.0	3.5				3.1	4.0	3.5	6.1	8.8	7.3
Trend 1998–2002 (per annum)				-4.9%	-1.9%	-3.1%				-5.4%	-0.3%	-2.4%

Table A81 – Diffuse non-Hodgkin's lymphomas (200)

South Australia Year – 2002	Incidence						Mortality					
	New cases			Rates			Deaths			Rates		
	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons
0–4	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
5–9	0	1	1	0.0	2.1	1.0	0	0	0	0.0	0.0	0.0
10–14	1	0	1	1.9	0.0	1.0	0	0	0	0.0	0.0	0.0
15–19	2	0	2	3.8	0.0	1.9	0	0	0	0.0	0.0	0.0
20–24	3	2	5	6.0	4.3	5.2	0	0	0	0.0	0.0	0.0
25–29	2	0	2	4.0	0.0	2.1	0	1	1	0.0	2.1	1.0
30–34	3	2	5	5.4	3.7	4.6	1	0	1	1.8	0.0	0.9
35–39	5	1	6	8.9	1.8	5.4	1	0	1	1.8	0.0	0.9
40–44	2	4	6	3.4	6.8	5.1	0	1	1	0.0	1.7	0.9
45–49	8	1	9	14.9	1.8	8.3	2	1	3	3.7	1.8	2.8
50–54	14	7	21	27.1	13.3	20.1	4	6	10	7.8	11.4	9.6
55–59	18	13	31	40.8	29.3	35.0	9	2	11	20.4	4.5	12.4
60–64	19	8	27	55.7	23.0	39.2	5	2	7	14.7	5.8	10.2
65–69	16	11	27	55.5	36.3	45.7	6	2	8	20.8	6.6	13.5
70–74	24	18	42	90.4	60.6	74.6	9	7	16	33.9	23.5	28.4
75–79	22	17	39	101.9	61.1	78.9	13	12	25	60.2	43.1	50.6
80–84	18	10	28	140.1	49.7	84.9	14	5	19	108.9	24.8	57.6
85+	10	9	19	124.6	49.2	72.2	8	5	13	99.7	27.4	49.4
Total/Crude rate	167	104	271	22.2	13.6	17.8	72	44	116	9.6	5.7	7.6
AS Rate (Aust)				21.5	11.3	15.9				9.4	4.6	6.6
Confidence Interval (95%)				18.3 – 24.8	9.1 – 13.5	14.0 – 17.8				7.2 – 11.6	3.2 – 6.0	5.4 – 7.8
AS Rate (World)				16.1	8.5	12.0				6.2	3.2	4.5
Confidence Interval (95%)				13.8 – 18.4	6.9 – 10.1	10.6 – 13.4				5.0 – 7.5	2.3 – 4.1	3.7 – 5.3
Risk				1 in 63	1 in 110	1 in 81				1 in 191	1 in 349	1 in 249
PYLL										513	335	848
% of all cancers	4.1	2.9	3.5				4.1	2.9	3.5	4.0	2.9	3.5
Trend 1998–2002 (per annum)				10.1%	-2.9%	3.1%				8.8%	-6.4%	0.3%

Note: All rates are expressed per 100,000 population. Age-standardised rates (AS Rate) use the Australian 2001 and New World Standard populations (see Methods)

Table A82 – Hodgkin's disease (201)

South Australia Year – 2002	Incidence						Mortality					
	New cases			Rates			Deaths			Rates		
	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons
0–4	1	0	1	2.2	0.0	1.1	0	0	0	0.0	0.0	0.0
5–9	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
10–14	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
15–19	3	2	5	5.7	4.0	4.8	0	0	0	0.0	0.0	0.0
20–24	1	1	2	2.0	2.1	2.1	0	0	0	0.0	0.0	0.0
25–29	3	1	4	6.1	2.1	4.1	1	0	1	2.0	0.0	1.0
30–34	2	1	3	3.6	1.9	2.7	1	0	1	1.8	0.0	0.9
35–39	2	0	2	3.6	0.0	1.8	1	0	1	1.8	0.0	0.9
40–44	2	3	5	3.4	5.1	4.3	0	0	0	0.0	0.0	0.0
45–49	0	1	1	0.0	1.8	0.9	0	0	0	0.0	0.0	0.0
50–54	1	1	2	1.9	1.9	1.9	0	0	0	0.0	0.0	0.0
55–59	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
60–64	3	0	3	8.8	0.0	4.4	0	0	0	0.0	0.0	0.0
65–69	1	4	5	3.5	13.2	8.5	0	0	0	0.0	0.0	0.0
70–74	2	1	3	7.5	3.4	5.3	1	1	2	3.8	3.4	3.6
75–79	1	0	1	4.6	0.0	2.0	0	0	0	0.0	0.0	0.0
80–84	0	1	1	0.0	5.0	3.0	0	1	1	0.0	5.0	3.0
85+	1	0	1	12.5	0.0	3.8	0	0	0	0.0	0.0	0.0
Total/Crude rate	23	16	39	3.1	2.1	2.6	4	2	6	0.5	0.3	0.4
AS Rate (Aust)				3.1	2.0	2.5				0.5	0.2	0.4
Confidence Interval (95%)				1.8 – 4.4	1.0 – 3.0	1.7 – 3.3				0.0 – 1.1	0.0 – 0.5	0.1 – 0.7
AS Rate (World)				2.9	1.9	2.4				0.5	0.1	0.3
Confidence Interval (95%)				1.7 – 4.1	0.9 – 2.8	1.6 – 3.2				0.0 – 1.0	0.0 – 0.3	0.1 – 0.6
Risk				1 in 415	1 in 565	1 in 477				1 in 2,131	1 in 5,945	1 in 3,126
PYLL										130	3	133
% of all cancers	0.6	0.4	0.5				0.6	0.4	0.5	0.2	0.1	0.2
Trend 1998–2002 (per annum)				5.5%	19.7%	11.5%				-8.8%	16.3%	-5.0%

Table A83 – Other lymphomas & histiocytic tissue tumours (202)

South Australia Year – 2002	Incidence						Mortality					
	New cases			Rates			Deaths			Rates		
	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons
0–4	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
5–9	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
10–14	0	1	1	0.0	2.0	1.0	0	0	0	0.0	0.0	0.0
15–19	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
20–24	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
25–29	1	0	1	2.0	0.0	1.0	0	0	0	0.0	0.0	0.0
30–34	1	2	3	1.8	3.7	2.7	0	0	0	0.0	0.0	0.0
35–39	2	1	3	3.6	1.8	2.7	1	0	1	1.8	0.0	0.9
40–44	1	1	2	1.7	1.7	1.7	3	0	3	5.1	0.0	2.6
45–49	4	6	10	7.5	11.0	9.2	0	0	0	0.0	0.0	0.0
50–54	8	8	16	15.5	15.2	15.3	1	2	3	1.9	3.8	2.9
55–59	9	12	21	20.4	27.0	23.7	1	2	3	2.3	4.5	3.4
60–64	3	6	9	8.8	17.3	13.1	1	2	3	2.9	5.8	4.4
65–69	4	6	10	13.9	19.8	16.9	2	4	6	6.9	13.2	10.2
70–74	6	4	10	22.6	13.5	17.8	2	3	5	7.5	10.1	8.9
75–79	11	4	15	51.0	14.4	30.3	2	3	5	9.3	10.8	10.1
80–84	0	3	3	0.0	14.9	9.1	1	2	3	7.8	9.9	9.1
85+	1	3	4	12.5	16.4	15.2	2	4	6	24.9	21.9	22.8
Total/Crude rate	51	57	108	6.8	7.4	7.1	16	22	38	2.1	2.9	2.5
AS Rate (Aust)				6.4	6.6	6.5				2.1	2.3	2.2
Confidence Interval (95%)				4.7 – 8.2	4.9 – 8.3	5.3 – 7.7				1.1 – 3.2	1.3 – 3.3	1.5 – 2.9
AS Rate (World)				5.0	5.4	5.2				1.5	1.6	1.6
Confidence Interval (95%)				3.7 – 6.2	4.0 – 6.8	4.2 – 6.1				0.8 – 2.2	0.9 – 2.3	1.1 – 2.1
Risk				1 in 205	1 in 178	1 in 191				1 in 701	1 in 536	1 in 604
PYLL										208	143	350
% of all cancers	1.2	1.6	1.4				1.2	1.6	1.4	0.9	1.5	1.1
Trend 1998–2002 (per annum)				10.1%	5.8%	7.4%				19.1%	9.1%	15.0%

Table A84 – Nodular lymphomas (202.0)

South Australia Year – 2002	Incidence						Mortality					
	New cases			Rates			Deaths			Rates		
	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons
0–4	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
5–9	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
10–14	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
15–19	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
20–24	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
25–29	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
30–34	0	2	2	0.0	3.7	1.8	0	0	0	0.0	0.0	0.0
35–39	2	1	3	3.6	1.8	2.7	1	0	1	1.8	0.0	0.9
40–44	1	1	2	1.7	1.7	1.7	3	0	3	5.1	0.0	2.6
45–49	4	3	7	7.5	5.5	6.5	0	0	0	0.0	0.0	0.0
50–54	7	8	15	13.6	15.2	14.4	1	2	3	1.9	3.8	2.9
55–59	9	8	17	20.4	18.0	19.2	1	2	3	2.3	4.5	3.4
60–64	2	3	5	5.9	8.6	7.3	1	2	3	2.9	5.8	4.4
65–69	3	6	9	10.4	19.8	15.2	1	4	5	3.5	13.2	8.5
70–74	3	4	7	11.3	13.5	12.4	2	3	5	7.5	10.1	8.9
75–79	6	3	9	27.8	10.8	18.2	1	3	4	4.6	10.8	8.1
80–84	0	3	3	0.0	14.9	9.1	1	2	3	7.8	9.9	9.1
85+	1	1	2	12.5	5.5	7.6	2	3	5	24.9	16.4	19.0
Total/Crude rate	38	43	81	5.1	5.6	5.3	14	21	35	1.9	2.7	2.3
AS Rate (Aust)				4.8	5.0	4.9				1.9	2.2	2.0
Confidence Interval (95%)				3.3 – 6.3	3.5 – 6.5	3.8 – 6.0				0.9 – 2.9	1.3 – 3.2	1.4 – 2.7
AS Rate (World)				3.8	4.0	3.9				1.3	1.6	1.5
Confidence Interval (95%)				2.7 – 4.9	2.9 – 5.2	3.1 – 4.7				0.7 – 2.0	0.9 – 2.3	1.0 – 1.9
Risk				1 in 270	1 in 228	1 in 247				1 in 798	1 in 536	1 in 637
PYLL										200	143	343
% of all cancers	0.9	1.2	1.1				0.9	1.2	1.1	0.8	1.4	1.1
Trend 1998–2002 (per annum)				14.5%	12.1%	13.1%				19.4%	14.2%	17.4%

Note: All rates are expressed per 100,000 population. Age-standardised rates (AS Rate) use the Australian 2001 and New World Standard populations (see Methods)

Table A85 – All lymphomas (200 201 202.0 202.1 202.2 202.8 202.9)

South Australia Year – 2002	Incidence						Mortality					
	New cases			Rates			Deaths			Rates		
	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons
0-4	1	0	1	2.2	0.0	1.1	0	0	0	0.0	0.0	0.0
5-9	0	1	1	0.0	2.1	1.0	0	0	0	0.0	0.0	0.0
10-14	1	1	2	1.9	2.0	2.0	0	0	0	0.0	0.0	0.0
15-19	5	2	7	9.4	4.0	6.8	0	0	0	0.0	0.0	0.0
20-24	4	3	7	8.1	6.4	7.2	0	0	0	0.0	0.0	0.0
25-29	6	1	7	12.1	2.1	7.2	1	1	2	2.0	2.1	2.1
30-34	6	5	11	10.9	9.3	10.1	2	0	2	3.6	0.0	1.8
35-39	9	2	11	16.1	3.6	9.9	3	0	3	5.4	0.0	2.7
40-44	5	8	13	8.6	13.6	11.1	3	1	4	5.1	1.7	3.4
45-49	12	8	20	22.4	14.6	18.5	2	1	3	3.7	1.8	2.8
50-54	23	16	39	44.6	30.3	37.4	5	8	13	9.7	15.2	12.5
55-59	27	25	52	61.2	56.3	58.7	10	4	14	22.7	9.0	15.8
60-64	25	14	39	73.4	40.3	56.7	6	4	10	17.6	11.5	14.5
65-69	21	21	42	72.9	69.3	71.1	8	6	14	27.8	19.8	23.7
70-74	32	23	55	120.5	77.4	97.7	12	11	23	45.2	37.0	40.9
75-79	34	21	55	157.5	75.4	111.3	15	15	30	69.5	53.9	60.7
80-84	18	14	32	140.1	69.6	97.0	15	8	23	116.7	39.8	69.7
85+	12	12	24	149.6	65.6	91.2	10	9	19	124.6	49.2	72.2
Total/Crude rate	241	177	418	32.1	23.1	27.5	92	68	160	12.2	8.9	10.5
AS Rate (Aust)				31.1	19.9	25.0				12.1	7.1	9.2
Confidence Interval (95%)				27.1 – 35.0	17.0 – 22.9	22.6 – 27.4				9.6 – 14.5	5.4 – 8.8	7.8 – 10.6
AS Rate (World)				24.0	15.8	19.6				8.3	4.9	6.4
Confidence Interval (95%)				21.1 – 26.9	13.4 – 18.1	17.8 – 21.4				6.8 – 9.8	3.8 – 6.1	5.5 – 7.3
Risk				1 in 44	1 in 61	1 in 51				1 in 141	1 in 204	1 in 167
PYLL										850	480	1,330
% of all cancers	5.9	4.9	5.4				5.9	4.9	5.4	5.1	4.5	4.8
Trend 1998-2002 (per annum)				9.7%	0.6%	4.8%				8.9%	-3.4%	2.3%

Table A86 – Other lymphomas (202.1 202.2 202.8 202.9)

South Australia Year – 2002	Incidence						Mortality					
	New cases			Rates			Deaths			Rates		
	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons
0-4	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
5-9	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
10-14	0	1	1	0.0	2.0	1.0	0	0	0	0.0	0.0	0.0
15-19	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
20-24	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
25-29	1	0	1	2.0	0.0	1.0	0	0	0	0.0	0.0	0.0
30-34	1	0	1	1.8	0.0	0.9	0	0	0	0.0	0.0	0.0
35-39	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
40-44	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
45-49	0	3	3	0.0	5.5	2.8	0	0	0	0.0	0.0	0.0
50-54	1	0	1	1.9	0.0	1.0	0	0	0	0.0	0.0	0.0
55-59	0	4	4	0.0	9.0	4.5	0	0	0	0.0	0.0	0.0
60-64	1	3	4	2.9	8.6	5.8	0	0	0	0.0	0.0	0.0
65-69	1	0	1	3.5	0.0	1.7	1	0	1	3.5	0.0	1.7
70-74	3	0	3	11.3	0.0	5.3	0	0	0	0.0	0.0	0.0
75-79	5	1	6	23.2	3.6	12.1	1	0	1	4.6	0.0	2.0
80-84	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
85+	0	2	2	0.0	10.9	7.6	0	1	1	0.0	5.5	3.8
Total/Crude rate	13	14	27	1.7	1.8	1.8	2	1	3	0.3	0.1	0.2
AS Rate (Aust)				1.7	1.6	1.6				0.2	0.1	0.2
Confidence Interval (95%)				0.8 – 2.5	0.8 – 2.5	1.0 – 2.2				0.0 – 0.6	0.0 – 0.2	0.0 – 0.4
AS Rate (World)				1.2	1.4	1.3				0.2	0.0	0.1
Confidence Interval (95%)				0.6 – 1.8	0.6 – 2.1	0.8 – 1.8				0.0 – 0.4	0.0 – 0.1	0.0 – 0.2
Risk				1 in 853	1 in 795	1 in 833				1 in 5,764	N/A	1 in 11,821
PYLL										8	0	8
% of all cancers	0.3	0.4	0.4				0.3	0.4	0.4	0.1	0.1	0.1
Trend 1998-2002 (per annum)				2.1%	-7.1%	-3.4%				16.3%	-14.9%	-0.5%

Table A87 – Tumors of histiocytic tissue (202.3 202.5 202.6)

South Australia Year – 2002	Incidence						Mortality					
	New cases			Rates			Deaths			Rates		
	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons
0-4	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
5-9	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
10-14	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
15-19	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
20-24	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
25-29	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
30-34	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
35-39	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
40-44	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
45-49	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
50-54	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
55-59	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
60-64	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
65-69	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
70-74	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
75-79	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
80-84	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
85+	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
Total/Crude rate	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
AS Rate (Aust)				0.0	0.0	0.0				0.0	0.0	0.0
Confidence Interval (95%)				0.0 – 0.0	0.0 – 0.0	0.0 – 0.0				0.0 – 0.0	0.0 – 0.0	0.0 – 0.0
AS Rate (World)				0.0	0.0	0.0				0.0	0.0	0.0
Confidence Interval (95%)				0.0 – 0.0	0.0 – 0.0	0.0 – 0.0				0.0 – 0.0	0.0 – 0.0	0.0 – 0.0
Risk				N/A	N/A	N/A				N/A	N/A	N/A
PYLL										0	0	0
% of all cancers	0.0	0.0	0.0				0.0	0.0	0.0	0.0	0.0	0.0
Trend 1998-2002 (per annum)				0.0%	0.0%	0.0%				0.0%	0.0%	0.0%

Note: All rates are expressed per 100,000 population. Age-standardised rates (AS Rate) use the Australian 2001 and New World Standard populations (see Methods)

Table A88 – Multiple myeloma & immunoproliferative neoplasms (203)

South Australia Year – 2002	Incidence						Mortality					
	New cases			Rates			Deaths			Rates		
	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons
0-4	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
5-9	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
10-14	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
15-19	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
20-24	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
25-29	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
30-34	1	0	1	1.8	0.0	0.9	0	0	0	0.0	0.0	0.0
35-39	2	0	2	3.6	0.0	1.8	0	0	0	0.0	0.0	0.0
40-44	1	0	1	1.7	0.0	0.9	0	0	0	0.0	0.0	0.0
45-49	3	1	4	5.6	1.8	3.7	1	0	1	1.9	0.0	0.9
50-54	4	5	9	7.8	9.5	8.6	3	2	5	5.8	3.8	4.8
55-59	5	1	6	11.3	2.3	6.8	4	1	5	9.1	2.3	5.6
60-64	9	5	14	26.4	14.4	20.4	2	1	3	5.9	2.9	4.4
65-69	7	6	13	24.3	19.8	22.0	5	3	8	17.4	9.9	13.5
70-74	15	7	22	56.5	23.5	39.1	6	6	12	22.6	20.2	21.3
75-79	17	6	23	78.8	21.6	46.5	12	9	21	55.6	32.3	42.5
80-84	6	8	14	46.7	39.8	42.5	2	10	12	15.6	49.7	36.4
85+	5	4	9	62.3	21.9	34.2	5	2	7	62.3	10.9	26.6
Total/Crude rate	75	43	118	10.0	5.6	7.8	40	34	74	5.3	4.4	4.9
AS Rate (Aust)				9.6	4.5	6.8				5.2	3.4	4.2
Confidence Interval (95%)				7.4 – 11.8	3.1 – 5.9	5.6 – 8.0				3.6 – 6.8	2.2 – 4.5	3.2 – 5.1
AS Rate (World)				6.7	3.2	4.8				3.4	2.2	2.7
Confidence Interval (95%)				5.3 – 8.1	2.2 – 4.1	4.0 – 5.6				2.5 – 4.4	1.5 – 2.8	2.2 – 3.3
Risk				1 in 144	1 in 281	1 in 193				1 in 320	1 in 513	1 in 396
PYLL										243	113	355
% of all cancers	1.8	1.2	1.5				1.8	1.2	1.5	2.2	2.3	2.2
Trend 1998-2002 (per annum)				-1.2%	-3.7%	-2.4%				-0.8%	-1.5%	-1.3%

Table A89 – All lymphoid leukaemias (204)

South Australia Year – 2002	Incidence						Mortality					
	New cases			Rates			Deaths			Rates		
	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons
0-4	5	0	5	10.8	0.0	5.5	0	0	0	0.0	0.0	0.0
5-9	3	3	6	6.0	6.3	6.1	0	0	0	0.0	0.0	0.0
10-14	1	0	1	1.9	0.0	1.0	0	0	0	0.0	0.0	0.0
15-19	2	0	2	3.8	0.0	1.9	0	1	1	0.0	2.0	1.0
20-24	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
25-29	0	1	1	0.0	2.1	1.0	0	1	1	0.0	2.1	1.0
30-34	0	0	0	0.0	0.0	0.0	1	1	2	1.8	1.9	1.8
35-39	1	0	1	1.8	0.0	0.9	0	0	0	0.0	0.0	0.0
40-44	1	2	3	1.7	3.4	2.6	1	1	2	1.7	1.7	1.7
45-49	1	3	4	1.9	5.5	3.7	0	0	0	0.0	0.0	0.0
50-54	3	2	5	5.8	3.8	4.8	0	2	2	0.0	3.8	1.9
55-59	6	3	9	13.6	6.8	10.2	0	0	0	0.0	0.0	0.0
60-64	8	5	13	23.5	14.4	18.9	2	0	2	5.9	0.0	2.9
65-69	6	5	11	20.8	16.5	18.6	2	0	2	6.9	0.0	3.4
70-74	12	8	20	45.2	26.9	35.5	5	3	8	18.8	10.1	14.2
75-79	16	4	20	74.1	14.4	40.5	6	2	8	27.8	7.2	16.2
80-84	8	8	16	62.3	39.8	48.5	3	1	4	23.3	5.0	12.1
85+	7	6	13	87.2	32.8	49.4	6	4	10	74.8	21.9	38.0
Total/Crude rate	80	50	130	10.6	6.5	8.6	26	16	42	3.5	2.1	2.8
AS Rate (Aust)				10.5	5.4	7.6				3.5	1.7	2.4
Confidence Interval (95%)				8.2 – 12.8	3.9 – 7.0	6.3 – 9.0				2.2 – 4.9	0.9 – 2.6	1.7 – 3.1
AS Rate (World)				7.9	4.2	5.9				2.2	1.3	1.6
Confidence Interval (95%)				6.2 – 9.7	2.9 – 5.4	4.8 – 7.0				1.4 – 2.9	0.6 – 2.0	1.1 – 2.1
Risk				1 in 147	1 in 234	1 in 181				1 in 569	1 in 930	1 in 716
PYLL										128	233	360
% of all cancers	2.0	1.4	1.7				2.0	1.4	1.7	1.4	1.1	1.3
Trend 1998-2002 (per annum)				3.8%	-4.8%	-0.3%				-2.3%	-8.2%	-5.5%

Table A90 – Acute lymphatic leukaemia (204.0)

South Australia Year – 2002	Incidence						Mortality					
	New cases			Rates			Deaths			Rates		
	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons
0-4	5	0	5	10.8	0.0	5.5	0	0	0	0.0	0.0	0.0
5-9	3	3	6	6.0	6.3	6.1	0	0	0	0.0	0.0	0.0
10-14	1	0	1	1.9	0.0	1.0	0	0	0	0.0	0.0	0.0
15-19	2	0	2	3.8	0.0	1.9	0	1	1	0.0	2.0	1.0
20-24	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
25-29	0	1	1	0.0	2.1	1.0	0	1	1	0.0	2.1	1.0
30-34	0	0	0	0.0	0.0	0.0	1	1	2	1.8	1.9	1.8
35-39	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
40-44	1	1	2	1.7	1.7	1.7	1	1	2	1.7	1.7	1.7
45-49	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
50-54	0	0	0	0.0	0.0	0.0	0	1	1	0.0	1.9	1.0
55-59	0	1	1	0.0	2.3	1.1	0	0	0	0.0	0.0	0.0
60-64	2	1	3	5.9	2.9	4.4	0	0	0	0.0	0.0	0.0
65-69	1	1	2	3.5	3.3	3.4	1	0	1	3.5	0.0	1.7
70-74	1	0	1	3.8	0.0	1.8	1	1	2	3.8	3.4	3.6
75-79	1	2	3	4.6	7.2	6.1	2	1	3	9.3	3.6	6.1
80-84	1	1	2	7.8	5.0	6.1	0	0	0	0.0	0.0	0.0
85+	1	0	1	12.5	0.0	3.8	2	0	2	24.9	0.0	7.6
Total/Crude rate	19	11	30	2.5	1.4	2.0	8	7	15	1.1	0.9	1.0
AS Rate (Aust)				2.6	1.3	1.9				1.1	0.9	0.9
Confidence Interval (95%)				1.4 – 3.7	0.5 – 2.2	1.2 – 2.6				0.3 – 1.9	0.2 – 1.6	0.5 – 1.4
AS Rate (World)				2.7	1.3	2.0				0.7	0.8	0.7
Confidence Interval (95%)				1.4 – 4.0	0.4 – 2.2	1.2 – 2.8				0.3 – 1.2	0.2 – 1.4	0.4 – 1.1
Risk				1 in 537	1 in 1,082	1 in 717				1 in 1,858	1 in 1,551	1 in 1,704
PYLL										85	205	290
% of all cancers	0.5	0.3	0.4				0.5	0.3	0.4	0.4	0.5	0.5
Trend 1998-2002 (per annum)				10.5%	-8.6%	0.4%				-3.0%	-14.0%	-9.8%

Note: All rates are expressed per 100,000 population. Age-standardised rates (AS Rate) use the Australian 2001 and New World Standard populations (see Methods)

Table A91 – Chronic lymphatic leukaemia (204.1)

South Australia Year – 2002	Incidence						Mortality					
	New cases			Rates			Deaths			Rates		
	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons
0–4	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
5–9	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
10–14	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
15–19	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
20–24	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
25–29	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
30–34	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
35–39	1	0	1	1.8	0.0	0.9	0	0	0	0.0	0.0	0.0
40–44	0	1	1	0.0	1.7	0.9	0	0	0	0.0	0.0	0.0
45–49	1	3	4	1.9	5.5	3.7	0	0	0	0.0	0.0	0.0
50–54	3	2	5	5.8	3.8	4.8	0	1	1	0.0	1.9	1.0
55–59	6	2	8	13.6	4.5	9.0	0	0	0	0.0	0.0	0.0
60–64	6	4	10	17.6	11.5	14.5	2	0	2	5.9	0.0	2.9
65–69	5	4	9	17.4	13.2	15.2	1	0	1	3.5	0.0	1.7
70–74	11	8	19	41.4	26.9	33.8	4	2	6	15.1	6.7	10.7
75–79	15	2	17	69.5	7.2	34.4	4	1	5	18.5	3.6	10.1
80–84	7	7	14	54.5	34.8	42.5	3	1	4	23.3	5.0	12.1
85+	6	6	12	74.8	32.8	45.6	4	4	8	49.9	21.9	30.4
Total/Crude rate	61	39	100	8.1	5.1	6.6	18	9	27	2.4	1.2	1.8
AS Rate (Aust)				7.9	4.1	5.7				2.4	0.8	1.5
Confidence Interval (95%)				5.9 – 9.9	2.8 – 5.4	4.6 – 6.8				1.3 – 3.6	0.3 – 1.4	0.9 – 2.1
AS Rate (World)				5.2	2.9	3.9				1.4	0.5	0.9
Confidence Interval (95%)				4.1 – 6.4	2.0 – 3.8	3.2 – 4.7				0.9 – 2.0	0.2 – 0.8	0.6 – 1.2
Risk				1 in 202	1 in 298	1 in 242				1 in 820	1 in 2,320	1 in 1,234
PYLL										43	28	70
% of all cancers	1.5	1.1	1.3				1.5	1.1	1.3	1.0	0.6	0.8
Trend 1998–2002 (per annum)				2.0%	-3.1%	-0.5%				-2.0%	-3.0%	-2.8%

Table A92 – All myeloid leukaemias (205)

South Australia Year – 2002	Incidence						Mortality					
	New cases			Rates			Deaths			Rates		
	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons
0–4	1	0	1	2.2	0.0	1.1	0	0	0	0.0	0.0	0.0
5–9	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
10–14	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
15–19	1	0	1	1.9	0.0	1.0	0	0	0	0.0	0.0	0.0
20–24	0	0	0	0.0	0.0	0.0	0	1	1	0.0	2.1	1.0
25–29	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
30–34	1	2	3	1.8	3.7	2.7	0	0	0	0.0	0.0	0.0
35–39	1	2	3	1.8	3.6	2.7	1	1	2	1.8	1.8	1.8
40–44	3	3	6	5.1	5.1	5.1	1	1	2	1.7	1.7	1.7
45–49	4	2	6	7.5	3.7	5.5	1	1	2	1.9	1.8	1.8
50–54	5	1	6	9.7	1.9	5.7	1	0	1	1.9	0.0	1.0
55–59	3	4	7	6.8	9.0	7.9	1	2	3	2.3	4.5	3.4
60–64	5	3	8	14.7	8.6	11.6	3	3	6	8.8	8.6	8.7
65–69	9	3	12	31.2	9.9	20.3	6	1	7	20.8	3.3	11.8
70–74	14	9	23	52.7	30.3	40.9	9	9	18	33.9	30.3	32.0
75–79	9	5	14	41.7	18.0	28.3	9	4	13	41.7	14.4	26.3
80–84	7	5	12	54.5	24.8	36.4	9	5	14	70.0	24.8	42.5
85+	7	7	14	87.2	38.3	53.2	7	8	15	87.2	43.8	57.0
Total/Crude rate	70	46	116	9.3	6.0	7.6	48	36	84	6.4	4.7	5.5
AS Rate (Aust)				9.2	4.9	6.8				6.4	3.7	4.7
Confidence Interval (95%)				7.0 – 11.3	3.5 – 6.4	5.5 – 8.0				4.6 – 8.2	2.4 – 4.9	3.7 – 5.8
AS Rate (World)				6.5	3.6	4.9				4.0	2.5	3.1
Confidence Interval (95%)				5.1 – 8.0	2.6 – 4.6	4.1 – 5.8				3.0 – 5.1	1.7 – 3.4	2.5 – 3.8
Risk				1 in 148	1 in 264	1 in 192				1 in 274	1 in 370	1 in 317
PYLL										243	253	495
% of all cancers	1.7	1.3	1.5				1.7	1.3	1.5	2.6	2.4	2.5
Trend 1998–2002 (per annum)				-2.8%	0.8%	-1.2%				-0.4%	-3.6%	-2.1%

Table A93 – Acute myeloid leukaemia (205.0)

South Australia Year – 2002	Incidence						Mortality					
	New cases			Rates			Deaths			Rates		
	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons
0–4	1	0	1	2.2	0.0	1.1	0	0	0	0.0	0.0	0.0
5–9	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
10–14	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
15–19	1	0	1	1.9	0.0	1.0	0	0	0	0.0	0.0	0.0
20–24	0	0	0	0.0	0.0	0.0	0	1	1	0.0	2.1	1.0
25–29	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
30–34	1	2	3	1.8	3.7	2.7	0	0	0	0.0	0.0	0.0
35–39	0	0	0	0.0	0.0	0.0	1	1	2	1.8	1.8	1.8
40–44	1	2	3	1.7	3.4	2.6	1	1	2	1.7	1.7	1.7
45–49	4	2	6	7.5	3.7	5.5	1	1	2	1.9	1.8	1.8
50–54	2	1	3	3.9	1.9	2.9	1	0	1	1.9	0.0	1.0
55–59	2	4	6	4.5	9.0	6.8	0	2	2	0.0	4.5	2.3
60–64	3	3	6	8.8	8.6	8.7	2	2	4	5.9	5.8	5.8
65–69	6	2	8	20.8	6.6	13.5	3	1	4	10.4	3.3	6.8
70–74	12	9	21	45.2	30.3	37.3	8	9	17	30.1	30.3	30.2
75–79	9	5	14	41.7	18.0	28.3	9	4	13	41.7	14.4	26.3
80–84	6	2	8	46.7	9.9	24.3	8	2	10	62.3	9.9	30.3
85+	6	7	13	74.8	38.3	49.4	6	8	14	74.8	43.8	53.2
Total/Crude rate	54	39	93	7.2	5.1	6.1	40	32	72	5.3	4.2	4.7
AS Rate (Aust)				7.1	4.2	5.4				5.3	3.3	4.1
Confidence Interval (95%)				5.2 – 9.0	2.8 – 5.5	4.3 – 6.5				3.7 – 7.0	2.1 – 4.4	3.1 – 5.0
AS Rate (World)				4.9	3.0	3.8				3.3	2.3	2.7
Confidence Interval (95%)				3.7 – 6.1	2.1 – 3.9	3.1 – 4.6				2.4 – 4.2	1.5 – 3.1	2.1 – 3.2
Risk				1 in 204	1 in 298	1 in 244				1 in 373	1 in 390	1 in 382
PYLL										188	240	428
% of all cancers	1.3	1.1	1.2				1.3	1.1	1.2	2.2	2.1	2.2
Trend 1998–2002 (per annum)				-2.3%	0.9%	-0.8%				-0.9%	-2.1%	-1.7%

Note: All rates are expressed per 100,000 population. Age-standardised rates (AS Rate) use the Australian 2001 and New World Standard populations (see Methods)

Table A94 – Chronic myeloid leukaemia (205.0)

South Australia Year – 2002	Incidence						Mortality					
	New cases			Rates			Deaths			Rates		
	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons
0–4	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
5–9	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
10–14	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
15–19	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
20–24	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
25–29	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
30–34	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
35–39	1	2	3	1.8	3.6	2.7	0	0	0	0.0	0.0	0.0
40–44	2	1	3	3.4	1.7	2.6	0	0	0	0.0	0.0	0.0
45–49	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
50–54	3	0	3	5.8	0.0	2.9	0	0	0	0.0	0.0	0.0
55–59	1	0	1	2.3	0.0	1.1	1	0	1	2.3	0.0	1.1
60–64	2	0	2	5.9	0.0	2.9	1	1	2	2.9	2.9	2.9
65–69	3	1	4	10.4	3.3	6.8	3	0	3	10.4	0.0	5.1
70–74	2	0	2	7.5	0.0	3.6	1	0	1	3.8	0.0	1.8
75–79	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
80–84	1	3	4	7.8	14.9	12.1	1	3	4	7.8	14.9	12.1
85+	1	0	1	12.5	0.0	3.8	1	0	1	12.5	0.0	3.8
Total/Crude rate	16	7	23	2.1	0.9	1.5	8	4	12	1.1	0.5	0.8
AS Rate (Aust)				2.1	0.8	1.4				1.0	0.4	0.7
Confidence Interval (95%)				1.1 – 3.1	0.2 – 1.4	0.8 – 2.0				0.3 – 1.8	0.0 – 0.7	0.3 – 1.1
AS Rate (World)				1.6	0.6	1.1				0.8	0.2	0.5
Confidence Interval (95%)				0.9 – 2.4	0.2 – 1.0	0.7 – 1.5				0.3 – 1.3	0.0 – 0.5	0.2 – 0.8
Risk				1 in 539	1 in 2,324	1 in 890				1 in 1,033	1 in 6,943	1 in 1,837
PYLL										55	13	68
% of all cancers	0.4	0.2	0.3				0.4	0.2	0.3	0.4	0.3	0.4
Trend 1998–2002 (per annum)				-4.0%	0.7%	-2.6%				2.9%	-10.2%	-4.2%

Table A95 – Chronic monocytic leukaemia (205.0)

South Australia Year – 2002	Incidence						Mortality					
	New cases			Rates			Deaths			Rates		
	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons
0–4	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
5–9	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
10–14	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
15–19	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
20–24	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
25–29	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
30–34	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
35–39	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
40–44	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
45–49	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
50–54	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
55–59	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
60–64	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
65–69	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
70–74	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
75–79	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
80–84	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
85+	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
Total/Crude rate	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
AS Rate (Aust)				0.0	0.0	0.0				0.0	0.0	0.0
Confidence Interval (95%)				0.0 – 0.0	0.0 – 0.0	0.0 – 0.0				0.0 – 0.0	0.0 – 0.0	0.0 – 0.0
AS Rate (World)				0.0	0.0	0.0				0.0	0.0	0.0
Confidence Interval (95%)				0.0 – 0.0	0.0 – 0.0	0.0 – 0.0				0.0 – 0.0	0.0 – 0.0	0.0 – 0.0
Risk				N/A	N/A	N/A				N/A	N/A	N/A
PYLL										0	0	0
% of all cancers	0.0	0.0	0.0				0.0	0.0	0.0	0.0	0.0	0.0
Trend 1998–2002 (per annum)				-25.0%	0.0%	-25.0%				-27.0%	0.0%	-25.5%

Table A96 – All leukaemias (204 205 206.1 207 208)

South Australia Year – 2002	Incidence						Mortality					
	New cases			Rates			Deaths			Rates		
	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons
0–4	6	0	6	13.0	0.0	6.6	0	0	0	0.0	0.0	0.0
5–9	3	3	6	6.0	6.3	6.1	0	0	0	0.0	0.0	0.0
10–14	1	0	1	1.9	0.0	1.0	0	0	0	0.0	0.0	0.0
15–19	3	0	3	5.7	0.0	2.9	0	1	1	0.0	2.0	1.0
20–24	0	0	0	0.0	0.0	0.0	0	1	1	0.0	2.1	1.0
25–29	0	1	1	0.0	2.1	1.0	0	1	1	0.0	2.1	1.0
30–34	1	2	3	1.8	3.7	2.7	1	1	2	1.8	1.9	1.8
35–39	2	2	4	3.6	3.6	3.6	1	1	2	1.8	1.8	1.8
40–44	5	5	10	8.6	8.5	8.5	2	2	4	3.4	3.4	3.4
45–49	5	5	10	9.3	9.1	9.2	1	2	3	1.9	3.7	2.8
50–54	8	4	12	15.5	7.6	11.5	1	2	3	1.9	3.8	2.9
55–59	11	7	18	24.9	15.8	20.3	1	2	3	2.3	4.5	3.4
60–64	15	9	24	44.0	25.9	34.9	6	3	9	17.6	8.6	13.1
65–69	18	8	26	62.5	26.4	44.0	9	2	11	31.2	6.6	18.6
70–74	30	18	48	113.0	60.6	85.3	14	13	27	52.7	43.7	48.0
75–79	29	9	38	134.4	32.3	76.9	18	6	24	83.4	21.6	48.6
80–84	17	13	30	132.3	64.6	91.0	14	7	21	108.9	34.8	63.7
85+	16	16	32	199.4	87.5	121.7	15	15	30	187.0	82.1	114.1
Total/Crude rate	170	102	272	22.6	13.3	17.9	83	59	142	11.0	7.7	9.4
AS Rate (Aust)				22.2	10.9	15.9				11.1	6.0	8.0
Confidence Interval (95%)				18.8 – 25.5	8.8 – 13.1	14.0 – 17.8				8.7 – 13.5	4.5 – 7.6	6.7 – 9.4
AS Rate (World)				16.2	8.2	11.9				6.9	4.2	5.3
Confidence Interval (95%)				13.8 – 18.6	6.5 – 9.8	10.4 – 13.3				5.6 – 8.2	3.1 – 5.4	4.5 – 6.1
Risk				1 in 65	1 in 118	1 in 85				1 in 175	1 in 238	1 in 203
PYLL										390	523	913
% of all cancers	4.2	2.8	3.5				4.2	2.8	3.5	4.6	3.9	4.3
Trend 1998–2002 (per annum)				0.9%	-3.1%	-0.9%				0.1%	-4.3%	-2.3%

Note: All rates are expressed per 100,000 population. Age-standardised rates (AS Rate) use the Australian 2001 and New World Standard populations (see Methods)

Table A97 – Monocytic leukaemias (chronic only) (206)

South Australia Year – 2002	Incidence						Mortality					
	New cases			Rates			Deaths			Rates		
	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons
0–4	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
5–9	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
10–14	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
15–19	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
20–24	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
25–29	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
30–34	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
35–39	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
40–44	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
45–49	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
50–54	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
55–59	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
60–64	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
65–69	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
70–74	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
75–79	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
80–84	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
85+	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
Total/Crude rate	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
AS Rate (Aust)				0.0	0.0	0.0				0.0	0.0	0.0
Confidence Interval (95%)				0.0 – 0.0	0.0 – 0.0	0.0 – 0.0				0.0 – 0.0	0.0 – 0.0	0.0 – 0.0
AS Rate (World)				0.0	0.0	0.0				0.0	0.0	0.0
Confidence Interval (95%)				0.0 – 0.0	0.0 – 0.0	0.0 – 0.0				0.0 – 0.0	0.0 – 0.0	0.0 – 0.0
Risk				N/A	N/A	N/A				N/A	N/A	N/A
PYLL										0	0	0
% of all cancers	0.0	0.0	0.0				0.0	0.0	0.0	0.0	0.0	0.0
Trend 1998–2002 (per annum)				-25.0%	0.0%	-25.0%				-27.0%	0.0%	-25.5%

Table A98 – Other specified leukaemias (207)

South Australia Year – 2002	Incidence						Mortality					
	New cases			Rates			Deaths			Rates		
	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons
0–4	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
5–9	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
10–14	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
15–19	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
20–24	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
25–29	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
30–34	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
35–39	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
40–44	1	0	1	1.7	0.0	0.9	0	0	0	0.0	0.0	0.0
45–49	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
50–54	0	1	1	0.0	1.9	1.0	0	0	0	0.0	1.0	0.0
55–59	2	0	2	4.5	0.0	2.3	0	0	0	0.0	0.0	0.0
60–64	2	1	3	5.9	2.9	4.4	1	0	1	2.9	0.0	1.5
65–69	2	0	2	6.9	0.0	3.4	0	0	0	0.0	0.0	0.0
70–74	3	1	4	11.3	3.4	7.1	0	1	1	0.0	3.4	1.8
75–79	3	0	3	13.9	0.0	6.1	2	0	2	9.3	0.0	4.0
80–84	2	0	2	15.6	0.0	6.1	2	1	3	15.6	5.0	9.1
85+	1	1	2	12.5	5.5	7.6	1	1	2	12.5	5.5	7.6
Total/Crude rate	16	4	20	2.1	0.5	1.3	6	3	9	0.8	0.4	0.6
AS Rate (Aust)				2.0	0.4	1.2				0.8	0.3	0.5
Confidence Interval (95%)				1.0 – 3.0	0.0 – 0.9	0.6 – 1.7				0.2 – 1.5	0.0 – 0.6	0.2 – 0.8
AS Rate (World)				1.4	0.3	0.8				0.5	0.2	0.3
Confidence Interval (95%)				0.8 – 2.1	0.0 – 0.6	0.5 – 1.2				0.2 – 0.8	0.0 – 0.3	0.1 – 0.4
Risk				1 in 659	1 in 2,458	1 in 1,057				1 in 6,817	1 in 5,945	1 in 6,192
PYLL										13	3	15
% of all cancers	0.4	0.1	0.3				0.4	0.1	0.3	0.3	0.2	0.3
Trend 1998–2002 (per annum)				1.8%	-14.2%	-4.8%				26.8%	15.2%	22.1%

Table A99 – Unspecified cell leukaemias (208)

South Australia Year – 2002	Incidence						Mortality					
	New cases			Rates			Deaths			Rates		
	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons
0–4	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
5–9	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
10–14	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
15–19	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
20–24	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
25–29	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
30–34	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
35–39	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
40–44	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
45–49	0	0	0	0.0	0.0	0.0	0	1	1	0.0	1.8	0.9
50–54	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
55–59	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
60–64	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
65–69	1	0	1	3.5	0.0	1.7	1	1	2	3.5	3.3	3.4
70–74	1	0	1	3.8	0.0	1.8	0	0	0	0.0	0.0	0.0
75–79	1	0	1	4.6	0.0	2.0	1	0	1	4.6	0.0	2.0
80–84	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
85+	1	2	3	12.5	10.9	11.4	1	2	3	12.5	10.9	11.4
Total/Crude rate	4	2	6	0.5	0.3	0.4	3	4	7	0.4	0.5	0.5
AS Rate (Aust)				0.5	0.1	0.3				0.4	0.4	0.4
Confidence Interval (95%)				0.0 – 1.1	0.0 – 0.4	0.1 – 0.6				0.0 – 0.9	0.0 – 0.8	0.1 – 0.7
AS Rate (World)				0.3	0.1	0.2				0.2	0.3	0.3
Confidence Interval (95%)				0.0 – 0.6	0.0 – 0.1	0.0 – 0.3				0.0 – 0.5	0.0 – 0.6	0.1 – 0.4
Risk				1 in 2,765	N/A	1 in 5,766				1 in 5,764	1 in 3,900	1 in 4,645
PYLL										8	35	43
% of all cancers	0.1	0.1	0.1				0.1	0.1	0.1	0.2	0.3	0.2
Trend 1998–2002 (per annum)				73.1%	-6.9%	15.8%				24.6%	7.4%	12.1%

Note: All rates are expressed per 100,000 population. Age-standardised rates (AS Rate) use the Australian 2001 and New World Standard populations (see Methods)

Table A100 – All cancers (140–208)

South Australia Year – 2002	Incidence						Mortality					
	New cases			Rates			Deaths			Rates		
	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons
0–4	9	6	15	19.4	13.5	16.5	2	0	2	4.3	0.0	2.2
5–9	7	7	14	13.9	14.6	14.2	1	2	3	2.0	4.2	3.1
10–14	6	6	12	11.6	12.3	11.9	2	1	3	3.9	2.0	3.0
15–19	22	12	34	41.6	23.8	32.9	3	3	6	5.7	5.9	5.8
20–24	16	17	33	32.2	36.1	34.1	1	1	2	2.0	2.1	2.1
25–29	29	32	61	58.6	67.4	62.9	5	4	9	10.1	8.4	9.3
30–34	44	68	112	79.8	126.0	102.6	10	5	15	18.1	9.3	13.7
35–39	74	81	155	132.3	146.1	139.1	15	19	34	26.8	34.3	30.5
40–44	99	154	253	169.9	261.5	215.9	17	31	48	29.2	52.6	41.0
45–49	147	207	354	273.9	378.3	326.6	33	54	87	61.5	98.7	80.3
50–54	244	343	587	472.8	649.9	562.3	83	96	179	160.8	181.9	171.5
55–59	350	353	703	793.6	794.7	794.1	131	99	230	297.0	222.9	259.8
60–64	431	353	784	1,264.6	1,016.9	1,139.6	143	110	253	419.6	316.9	367.8
65–69	527	382	909	1,828.8	1,261.2	1,538.0	219	150	369	760.0	495.2	624.3
70–74	735	411	1,146	2,767.6	1,382.7	2,036.2	288	173	461	1,084.5	582.0	819.1
75–79	652	436	1,088	3,020.8	1,566.0	2,201.3	356	229	585	1,649.4	822.5	1,183.6
80–84	410	363	773	3,190.7	1,803.7	2,344.2	269	235	504	2,093.4	1,167.7	1,528.4
85+	289	386	675	3,602.1	2,111.7	2,566.3	243	291	534	3,028.8	1,592.0	2,030.3
Total/Crude rate	4,091	3,617	7,708	544.5	471.3	507.5	1,821	1,503	3,324	242.4	195.9	218.9
AS Rate (Aust)				526.7	398.0	451.9				239.4	153.2	188.7
Confidence Interval (95%)				510.5 – 542.9	384.8 – 411.0	441.7 – 462.0				228.3 – 250.4	145.3 – 161.1	182.3 – 195.2
AS Rate (World)				378.2	303.8	335.5				157.5	106.3	128.1
Confidence Interval (95%)				367.3 – 389.1	293.9 – 313.7	328.2 – 342.8				151.0 – 164.1	100.9 – 111.7	124.0 – 132.3
Risk				1 in 3	1 in 4	1 in 3				1 in 7	1 in 10	1 in 9
PYLL										11,558	10,855	22,413
% of all cancers	100.0	100.0	100.0				100.0	100.0	100.0	100.0	100.0	100.0
Trend 1998–2002 (per annum)				0.0%	0.0%	0.0%				0.0%	0.0%	0.0%

Table A101 – Cancer of the Head and Neck (141–149)

South Australia Year – 2002	Incidence						Mortality					
	New cases			Rates			Deaths			Rates		
	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons	Males	Females	Persons
0–4	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
5–9	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
10–14	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
15–19	1	0	1	1.9	0.0	1.0	0	0	0	0.0	0.0	0.0
20–24	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
25–29	0	0	0	0.0	0.0	0.0	0	0	0	0.0	0.0	0.0
30–34	1	1	2	1.8	1.9	1.8	0	0	0	0.0	0.0	0.0
35–39	7	1	8	12.5	1.8	7.2	0	0	0	0.0	0.0	0.0
40–44	6	0	6	10.3	0.0	5.1	1	1	2	1.7	1.7	1.7
45–49	7	5	12	13.0	9.1	11.1	4	1	5	7.5	1.8	4.6
50–54	13	4	17	25.2	7.6	16.3	4	0	4	7.8	0.0	3.8
55–59	10	4	14	22.7	9.0	15.8	6	0	6	13.6	0.0	6.8
60–64	12	1	13	35.2	2.9	18.9	7	1	8	20.5	2.9	11.6
65–69	9	2	11	31.2	6.6	18.6	3	0	3	10.4	0.0	5.1
70–74	14	6	20	52.7	20.2	35.5	4	1	5	15.1	3.4	8.9
75–79	11	7	18	51.0	25.1	36.4	4	1	5	18.5	3.6	10.1
80–84	4	2	6	31.1	9.9	18.2	5	2	7	38.9	9.9	21.2
85+	2	5	7	24.9	27.4	26.6	4	4	8	49.9	21.9	30.4
Total/Crude rate	97	38	135	12.9	5.0	8.9	42	11	53	5.6	1.4	3.5
AS Rate (Aust)				12.3	4.1	8.1				5.4	1.1	3.1
Confidence Interval (95%)				9.9 – 14.8	2.8 – 5.5	6.7 – 9.5				3.8 – 7.1	0.4 – 1.7	2.2 – 3.9
AS Rate (World)				9.7	3.0	6.3				3.9	0.7	2.2
Confidence Interval (95%)				7.8 – 11.5	2.1 – 4.0	5.2 – 7.3				2.8 – 5.1	0.3 – 1.1	1.6 – 2.8
Risk				1 in 97	1 in 339	1 in 153				1 in 262	1 in 2,047	1 in 471
PYLL										458	75	533
% of all cancers	2.4	1.1	1.8				2.4	1.1	1.8	2.3	0.7	1.6
Trend 1998–2002 (per annum)												

Note: All rates are expressed per 100,000 population. Age-standardised rates (AS Rate) use the Australian 2001 and New World Standard populations (see Methods)



Department
of Health

Cancer in SA 2002

This publication draws together the most recent cancer incidence and mortality data for South Australia. It projects these data forward to 2005 and examines the trends and patterns of cancer in the community. It highlights the impacts of smoking and alcohol and highlights thirteen cancers for a discussion of their risk factors and management.